

ONE DIMENSIONAL STRUCTURES IN O-MINIMAL THEORIES

ASSAF HASSON*, ALF ONSHUUS, AND YA'ACOV PETERZIL

ABSTRACT. We prove the Zil'ber Trichotomy Principle for all 1-dimensional structures which are definable in o-minimal ones.

On the level of types, we prove the following: Let $\mathcal{M}$ be an o-minimal structure and let $\mathcal{N}$ be any structure definable in $\mathcal{M}$. Assume that p is an $\mathcal{N}$ -type which is 1-dimensional in the sense of $\mathcal{M}$. Then, essentially, $\mathcal{N}$ induces on p either a trivial $acl_{\mathcal{N}}$ -geometry, or the structure of a pure (possibly ordered) $\mathcal{N}$ -definable vector space over a division ring, or an $\mathcal{N}$ -definable o-minimal expansion of a real closed field.

In particular, if $\mathcal{N}$ is stable and 1-dimensional then the acl geometry is necessarily locally modular.

Along the way, we develop a fine intersection theory for definable curves in o-minimal structures.

The present work is a first step in an attempt to classify structures interpretable in o-minimal theories. Such structures may be stable (e.g., algebraically closed fields of characteristic zero, compact complex manifolds) or unstable (e.g., expansions of real closed fields, ordered vector spaces). The ultimate goal would be to treat in some uniform manner all of these examples by exploiting the ambient o-minimal environment.

Definition 0.1. A structure $\mathcal{N}$ is *definable* in an o-minimal structure $\mathcal{M}$ if it is interpretable in the real sorts of $\mathcal{M}$ (i.e. not in $\mathcal{M}^{eq}$). More precisely, the universe of $\mathcal{N}$, call it N , is an $\mathcal{M}$ -definable set and its atomic relations are certain $\mathcal{M}$ -definable subsets of N^k , $k \in \mathbb{N}$. We say that $\mathcal{N}$ is *definable of dimension k in $\mathcal{M}$* if k is the minimal possible dimension of such an $\mathcal{M}$ -definable set N .

If $\mathcal{N}$ is definable in an o-minimal structure $\mathcal{M}$, quite a few good properties are induced on $\mathcal{N}$. Here is a partial list:

- (1) $\mathcal{N}$ has the non-independence property, NIP.
- (2) $\mathcal{N}$ is super-rosy of finite $\mathfrak{p}$ -rank, [16]
- (3) There is a bound on uniformly definable families of finite sets in $\mathcal{M}$ and therefore also in $\mathcal{N}$. Said differently, $\mathcal{N}$ eliminates the $\exists^\infty$ quantifier.
- (4) If N is a $\mathcal{N}$ -minimal (i.e. N has no $\mathcal{N}$ -definable infinite subsets of smaller $\mathcal{M}$ -dimension) then $acl_{\mathcal{N}}$ (the algebraic closure in the sense of $\mathcal{N}$) satisfies the exchange property (see [19]). Combined with (3), this means that $\mathcal{N}$ is a geometric structure (see [14] for more on geometric structures).
- (5) If $\mathcal{N}$ is stable then it has finite U -rank, with $U(\mathcal{N}) \leq \dim_{\mathcal{M}}(N)$, [7].

Let us consider the following setting: *Fix $\mathcal{M}$, a sufficiently saturated o-minimal structure with a dense underlying order*, and assume that $\mathcal{N}$ is a geometric structure definable in $\mathcal{M}$.

*Supported by the EPSRC grant no. EP C52800X 1.

Our main interest is to extract algebraic information from the geometric structure of $\mathcal{N}$. The idea is to follow Zilber's division of geometric structures $\mathcal{N}$ into three types:

Degenerate: $\text{acl}_{\mathcal{N}}(A) = \bigcup \{\text{acl}_{\mathcal{N}}(a) : a \in A\}$ for all $A \subset \mathcal{N}$.

Linear: Every normal definable family of plane curves in $\mathcal{N}$ is 1-dimensional (in the sense of $\mathcal{N}$), but $\mathcal{N}$ is non-degenerate.

Rich: There exists a normal 2-dimensional (in the sense of $\mathcal{N}$) definable family of plane curves.

Clearly, degenerate geometries do not allow the existence of infinite definable groups, and indeed by compactness not even the existence of type definable ones. Therefore there is no point looking for algebraic data in degenerate structures. But the Linear/Rich dichotomy has been subject to much research. The underlying thesis, sometimes known as Zilber's Principle, is that linear geometries should arise from pure linear structures (e.g a vector space with no additional structure) and that in contexts of "topological flavour" rich combinatorial geometries arise from the geometry of interpretable fields.

In the stable case, though not in general true (see [13]), Zilber's Principle has been first proved in [15] for strongly minimal Zariski geometries, and extended later to several related contexts. In the o-minimal world a local version of the principle was proved in [20].

A natural conjecture to make in this context is:

Conjecture *If $\mathcal{N}$ is a geometric structure definable in an o-minimal $\mathcal{M}$ then Zilber's Principle holds for $\mathcal{N}$. In particular, if $\mathcal{N}$ is rich then it interprets a field.*

Note that the above conjecture, if true, will cover some remaining open cases of Zilber's original conjecture, for strongly minimal structures interpretable in algebraically closed fields of characteristic zero (See Rabinovich's work [24] on the main cases of this question).

Our present work can be seen as a proof of the above conjecture in the case where $\mathcal{N}$ is definable in $\mathcal{M}$ and $\dim_{\mathcal{M}}(N) = 1$ (see [8] and [21] for the investigation of certain cases of definable, stable, two-dimensional structures). First note that if $\mathcal{N}$ is 1-dimensional and definable in $\mathcal{M}$ then every $\mathcal{N}$ -definable subset of N is either finite or has dimension one. Thus N itself is an $\mathcal{N}$ -minimal set and therefore a geometric structure. The main bulk of our work goes into proving that if $\mathcal{N}$ is one-dimensional, definable in $\mathcal{M}$, and stable then it cannot be rich. Let us see, in view of Zilber's Principle above, why we should indeed expect this to be true:

Assume that $\mathcal{N}$ is rich, then a field K should be interpretable in $\mathcal{N}$ (if we believe the above principle). Because $\mathcal{N}$ is superstable the field K must be algebraically closed, and because it is interpretable in an o-minimal theory it has to be of characteristic zero. Moreover, if $\mathcal{M}$ expands a real closed field R then K is definably isomorphic to the algebraic closure of R and in particular its o-minimal dimension is two (for both statements see [18]). It follows ([21]) that K , with all its $\mathcal{N}$ -induced structure, is a pure field and therefore strongly minimal. However, because K is interpretable in the geometric structure $\mathcal{N}$, its strong minimality implies that $\dim_{\mathcal{M}}(K) = \dim_{\mathcal{M}}(N) = 1$, a contradiction. (The assumption that $\mathcal{M}$ expands a real closed field can actually be avoided).

The same argument shows, assuming that the conjecture is true: If $\mathcal{N}$ is a rich strongly minimal structure which is interpretable in an o-minimal one then necessarily $\dim_{\mathcal{M}}(N) = 2$. Interestingly, the remaining open cases in proving Zilber's conjecture for strongly minimal structures in algebraically closed fields are exactly those in which the underlying universe of $\mathcal{N}$ is of *algebraic dimension* bigger than one.

Returning to our present work, after proving that a stable one-dimensional structure definable in $\mathcal{M}$ cannot be rich, the analysis of the linear stable case reduces to classical structure theorems on 1-based theories of U -rank 1. In the unstable case, we rely on results from [9] and [20] to finish our analysis. Taken together, our main theorem is:

Theorem 1. *Let $\mathcal{N}$ be a 1-dimensional structure definable in an o-minimal one. Then one of the following hold:*

- (1) $\mathcal{N}$ is degenerate.
- (2) $\mathcal{N}$ is linear: There exists $X \subseteq N^{eq}$ such that either X is strongly minimal and locally modular (whose $\mathcal{N}$ -definable structure arises from a definable vector space) or X is an o-minimal group interval, whose $\mathcal{N}$ -definable structure arises from an ordered vector space.
- (3) $\mathcal{N}$ is rich and interprets a real closed field.

Because of the flexibility in patching together any two o-minimal structures into a new one, it is clear that on the global level we cannot hope for sharper results. To get the more informative statement, our theorem will be formulated in terms of certain types in an arbitrary $\mathcal{M}$ -definable structure $\mathcal{N}$. We say that an $\mathcal{N}$ -type p is *$\mathcal{M}$ -1-dimensional* if it is contained in some $\mathcal{N}$ -definable set whose $\mathcal{M}$ -dimension is one. Because $\mathcal{N}$ has finite $\mathfrak{p}$ -rank, a natural first step in any analysis of such an $\mathcal{N}$ would be a classification of its $\mathfrak{p}$ -rank 1 types. Here we complete this classification for the case when $\mathfrak{p}$ -rank minimal types are $\mathcal{M}$ -1-dimensional. The work of [9] suggests that for unstable types this classification is not far from being sufficient. For a non-algebraic type $p \in S_1(N)$ say that:

Trivial: p is trivial if for every finite set of parameters $A \subseteq N$, and $a_1, \dots, a_n, b \models p$, if $b \in \text{acl}_{\mathcal{N}}(Aa_1 \dots a_n)$ then $b \in \bigcup_{i=1}^n \text{acl}_{\mathcal{N}}(Aa_i)$.

Rich: p is rich if there exist a finite set A , an element $b \models p$ with $\dim_{\mathcal{N}}(b/AN) = 1$ and an (almost) normal family $\mathcal{F}$ of plane curves, $\mathcal{N}$ -definable over the set NA , such that $\{f \in \mathcal{F} : \langle b, b \rangle \in f\}$ is infinite.

Linear: p is linear if p is not trivial and not rich.

Note that the notion of richness is definable in the following sense: If p is rich, witnessed by b and a family of curves $\mathcal{F}_{\bar{a}}$ (over parameters in $\bar{a}$) then, because $\dim_{\mathcal{N}}(b/Na) = 1$, there exists an $\bar{a}$ -definable infinite set $X_{\bar{a}}$ containing b such that whenever $b' \in X_{\bar{a}}$ the pair $\langle b', b' \rangle$ belongs to infinitely many curves from $\mathcal{F}_{\bar{a}}$. We can now find a formula $\theta \in p$ such that for all $b' \models \theta$ there exist a tuple $\bar{a}'$, $\mathcal{F}_{\bar{a}'}$ and $X_{\bar{a}'}$ with the same properties. We will say that θ isolates the richness of p .

The local version of Theorem 1 is then:

Theorem 2. *Assume that $\mathcal{N}$ is a definable structure in an o-minimal $\mathcal{M}$ and that p is a complete 1- $\mathcal{M}$ -dimensional $\mathcal{N}$ -type over a model $\mathcal{N}_0 \subseteq \mathcal{N}$. Then:*

- (1) p is trivial (with respect to $\text{acl}_{\mathcal{N}}$).

Or, there exists an $\mathcal{N}$ -definable equivalence relation E with finite classes such that one of the following holds:

- (2) *p is linear, in which case either*
 - (a) *p/E is a generic type of a strongly minimal $\mathcal{N}$ -definable one-dimensional group G , and the structure which $\mathcal{N}$ induces on G is locally modular. In particular, p is strongly minimal. Or,*
 - (b) *for every $a \models p$ there exists an $\mathcal{N}$ -definable ordered group-interval I containing a/E . The structure which $\mathcal{N}$ induces on I is a reduct of an ordered vector space over an ordered division ring.*
- (3) *p is rich: For every $a \models p$ there exists an $\mathcal{N}$ -definable real closed field R containing a and the structure which $\mathcal{N}$ induces on R is o-minimal.*

The above statement gives a more precise meaning to Theorem 1 and clearly strengthen it.

The results of this paper are true for $\mathcal{N}$ *definable* in arbitrary dense o-minimal structures (and by elimination of imaginaries, we may replace "definable" with "interpretable", if $\mathcal{M}$ expands an o-minimal group). However, in order to keep the exposition cleaner, throughout the main part of the work, we will assume that in fact $\mathcal{M}$ expands an o-minimal field. In Appendix B we prove the results to arbitrary o-minimal structures.

The structure of the paper Section 1 is quite technical. In it we develop a fine theory of tangency and transversality for curves in definable families. An important ingredient in the whole argument turns out to be the theory of limit sets as developed by v.d. Dries in [5] and we review it here and in Appendix A. In Section 2 we prove that every stable definable 1-dimensional structure is necessarily 1-based. In Section 3 we prove the main theorem, for structures and for types. As mentioned above, in Appendix B we generalise the result from expansions of real closed fields to arbitrary o-minimal structures.

1. SOME ELEMENTS OF INTERSECTION THEORY

In this section we develop elements of intersection theory which will be used in the proof of Theorem 1. Some parts of this theory were already developed in [20] and we will use several results and proofs from that paper. However, the setting there allowed us to “trim” the original family of curves as the proof progressed, by working locally in regions where the original family is “well-behaved”. In the current setting this is impossible (because the ordering is not assumed to be definable in $\mathcal{N}$) and therefore one needs to develop a finer intersection theory. We mainly concentrate on counting the number of intersection points of plane curves.

Although the treatment of the strongly minimal case will focus on reaching a contradiction (from the assumption that the structure is one dimensional and locally modular) the results proved in this section are true in any o-minimal structure with a family of curves as given below. Some of the results are stated under the assumption that the o-minimal structure expands a field. In Appendix B we will show how to avoid this assumption.

1.1. Intersections of two curves. We use the term *curve* to denote any $\mathcal{M}$ -definable one-dimensional set. A *plane curve* is a curve in M^2 .

Definition 1.1. Given an $\mathcal{M}$ -definable plane curve and $p \in X$, we say that X is C^0 at p if X is locally $\mathcal{M}$ -homeomorphic to an open interval (we also say that p is a C^0 -point of X).

It is not difficult to see that in this case X divides M^2 , sufficiently close to p , into two definably connected components, W_1, W_2 . Namely, for every sufficiently small rectangular neighbourhood W of p , the set $W \setminus X$ has two definably connected components, one having the same germ as W_1 at p and the other as W_2 .

The following definitions come from 2.13 of [20]:

Definition 1.2. If X and Y are two $\mathcal{M}$ -definable plane curves and p is a C^0 -point on both, we say that X and Y touch each other at p (see Figure 1 below) if there is some rectangular neighbourhood U of p such that $U \setminus X$ has two components W_1, W_2 and either $Y \cap U \cap W_1 = \emptyset$ or $Y \cap U \cap W_2 = \emptyset$.




Figure 1:
 X touches Y at p




Figure 2:
 $X \prec_p^+ Y, Y \prec_p^- X$

It is easy to see that “touching” is a symmetric relation. If $\mathcal{M}$ expands a real closed field and X and Y are C^1 -curves then this notion of tangency is stronger than the two curves having the same tangent space at p .

Definition 1.3. Let X and Y be two definable plane curves. Assume that $p = \langle x_0, y_0 \rangle \in X \cap Y$ and that X and Y are graphs of functions, $f(x)$ and $g(x)$, respectively, near p_0 . We say that $X \prec_p^+ Y$ ($X \prec_p^- Y$) if there is $x_1 > x_0$ such that $f(x) \leq g(x)$ ($f(x) < g(x)$) for all x in the interval (x_0, x_1) .

We define $X \prec_p^- Y$ ($X \prec_p^+ Y$) if there exists $x_1 < x_0$ such that for all $x \in (x_1, x_0)$, we have $f(x) \leq g(x)$ ($f(x) < g(x)$). We write $X \preceq_p Y$ ($X \prec_p Y$) if $Y \preceq_p^- X$ ($Y \prec_p^- X$) and $X \preceq_p^+ Y$ ($X \prec_p^+ Y$).

If X and Y are graphs of C^1 functions $f(x), g(x)$, with respect to a definable real closed field, and $f(x_0) = g(x_0)$, $f'(x_0) < g'(x_0)$ then we have $X \prec_p Y$ (see Figure 2)

Notice that, in the setting above, X and Y touch each other at p if and only one of the following holds:

- (i) $X \preceq_p^- Y$ and $X \prec_p^+ Y$ (namely, Y touches X from “below”) , or
- (ii) $Y \preceq_p^- X$ and $Y \prec_p^+ X$ (Y touches X from “above”).

1.2. Normal families of curves.

Definition 1.4. A definable family of curves $\mathcal{F} = \{C_q : q \in Q\}$ in M^n is called normal if for every $q_1, q_2 \in Q$, the curves C_{q_1} and C_{q_2} intersect in at most finitely many points. We say that a normal $\mathcal{F}$ has dimension k if $\dim Q = k$.

A family $\mathcal{F}$ as above is called *almost normal* if for every $q_1 \in Q$ there are at most finitely many $q_2 \in Q$ such that $C_{q_1} \cap C_{q_2}$ is infinite. The dimension of $\mathcal{F}$ is by definition $\dim(Q)$.

Note that if $\mathcal{F}$ is almost normal and $q \in Q$ is generic, there exists a neighbourhood $U \ni q$ such that $C_q \cap C_{q'}$ is finite for all $q' \in U$. If we take U definable over generic parameters, then we get by the genericity of q , an open $V \ni q$ such that for all $q_1, q_2 \in V$, $C_{q_1} \cap C_{q_2}$ is finite. Therefore, $\mathcal{F}_V := \{C_q : q \in V\}$ is normal. Since the results of this section are all local in nature, using the above observation, they are all true for almost normal families (with the same proofs). For the sake of clarity, the statements are only given for normal families.

Lemma 1.5. *Let $\mathcal{F} = \{C_q : q \in Q\}$ be a definable, k -dimensional normal family of curves in $P \subseteq M^n$, with $\dim P = 2$ and $k > 1$. Assume that $\mathcal{F}$ is definable over A .*

(i) *For every $\langle p, q \rangle \in P \times Q$, if $p \in C_q$ and $\dim(p, q/A) = k + 1$ then p and q are generic over A in P and Q , respectively, and $\dim(p/qA) = 1$.*

(ii) *If $P \subseteq M^2$, $p = \langle x_0, y_0 \rangle$ and $\dim_{\mathcal{M}}(p, q) = k + 1$ then there exist open intervals I and J , around x_0 and y_0 , respectively, such that $C_q \cap I \times J$ is the graph of a continuous, strictly monotone function.*

Proof. (i) Because for every $q \in Q$ we have $\dim(C_q) = 1$, it follows that $\dim(q/A) = k$, and hence $\dim(p/qA) = 1$. Because the intersection of any two C_q is finite we have $\dim(p/A) = 2$.

(ii) Because p is generic in C_q , the curve C_q is the graph of a continuous function f_q near p (either in x or in y). If this function were locally constant, say as a function of x , then, by the genericity of q we would get a k -dimensional set of f_q locally constant near x_0 . Because $k > 1$ we would get an infinite set of q' such that $f_{q'}$ all agree on a whole interval, contradicting normality. $\square$

When $\mathcal{F}$ is a fixed family of curves parameterised by Q then we sometimes write $q_1 \preceq_p q_2$ instead of $C_{q_1} \preceq_p C_{q_2}$.

1.3. Nice families of curves.

Lemma 1.6. *Let $\mathcal{F} = \{C_q : q \in Q\}$ be a $\emptyset$ -definable two-dimensional normal family of curves in $P \subseteq M^2$. Assume that $p_0 \in C_{q_0}$ and that $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$. Then there exist a neighbourhood $U \times W \subseteq M^2 \times Q$ of $\langle p_0, q_0 \rangle$ such that for all $q_1, q_2 \in W$ the curves C_{q_1} and C_{q_2} intersect at most once in U and they do not touch each other at their point of intersection.*

Proof. Assume that p_0 and q_0 are as above. Then, by possibly shrinking P and Q we may assume that for every $q \in Q$ the curve $C_q \subseteq P$ is the graph of a continuous partial function f_q from M into M and that f_q vary continuously with q .

Claim 1 For all $q \neq q_0$ sufficiently close to q_0 the curve C_q does not touch C_{q_0} at p_0 .

Indeed, if the claim fails then there is a 1-dimensional set $Q' \subseteq Q$ containing q_0 and definable over generic parameters, such that $C_{q'}$ touches C_{q_0} at p_0 for every $q' \in Q'$. Because q_0 is generic, by standard dimension considerations, we can obtain a definable, one-dimensional set Q_{p_0} such that for any $q_1, q_2 \in Q_{p_0}$, the curves C_{q_1}, C_{q_2} , touch each other at p_0 . Now, by varying p_0 , we may find an open set U containing p_0 and a relatively open $Q_1 \subseteq Q$ such that for every $p \in U$ and every $q_1, q_2 \in Q_1$,

if $p \in C_{q_1} \cap C_{q_2}$ then C_{q_1} and C_{q_2} touch each other at p . We may assume that U and Q_1 are 0-definable (by choosing them to be definable over sufficiently generic and independent parameters). Let us show that this contradicts the fact that Q is two-dimensional and $\mathcal{F}$ is normal:

Let $p_0 = \langle x_0, y_0 \rangle$ and $Q_0 = \{q \in Q_1 : f_q(x_0) = y_0\}$. Choose $x_1 > x_0$ generic (over all mentioned parameters) and $y_1 = f_{q_0}(x_1)$ such that $\langle x_1, y_1 \rangle \in U$. Since y_1 is generic in M over $p_0 q_0$, there are $q_1, q_2 \in Q_{p_0}$ such that $f_{q_1}(x_1) < y_1 < f_{q_2}(x_1)$. If we now take any $q \in Q_1$ such that $f_q(x_0) \neq y_0$ then necessarily $f_q(x_1) \neq y_1$, for otherwise the graph of f_q will have to cross (and therefore not touch) either C_{q_1} or C_{q_2} in the interval (x_0, x_1) . By the normality of $\mathcal{F}$ and the genericity of $(x_0, y_0), (x_1, y_1)$ there are at most finitely many $q \in Q_1$ such that both points are in C_q . So there are at most finitely many $q \in Q_1$ such that $\langle x_1, y_1 \rangle \in C_{q_1}$, in contradiction to the fact that $\mathcal{F}$ is two-dimensional. End of Claim 1.

Claim 2 There is an open $U \subseteq P$ and an open $W \subseteq Q$ such that for every $q_1 \neq q_2 \in U$ and for every $p \in C_{q_1} \cap C_{q_2} \cap W$, the curves C_{q_1} and C_{q_2} do not touch each other at p .

Indeed, by Claim 1 there is a relatively open $W' \subseteq Q$ such that for every $q \neq q_0$ in W' , if $p_0 \in C_q \cap C_{q_0}$ then C_q and C_{q_0} do not touch each other at p_0 . By choosing the parameters defining W' sufficiently independent, we may assume that W' is 0-definable. By the genericity of q_0 in Q_{p_0} , we may find an open $W \subseteq Q$, $q_0 \in W \subseteq W'$, such that for every $q', q'' \in W$, if $p_0 \in C_{q'} \cap C_{q''}$ then the two curves do not touch each other at p_0 . Again, we assume that W is 0-definable. Finally, we may use the genericity of p_0 to obtain the desired U . End of Claim 2.

The rest of the proof is extracted from Lemma 4.3 in [20], so we only sketch the argument.

We first may assume that $Q \subseteq M^2$. By Claim 2, we may assume that for all $p \in U$ and $q_1, q_2 \in Q$, if C_{q_1}, C_{q_2} intersect in $p \in U$ then either $q_1 \preceq_p q_2$ or $q_2 \preceq_p q_1$.

Using Claim 2 and o-minimal-type dimension arguments, we may find open definable neighbourhoods W of q_0 and U of p_0 such that for every $p \in U$ and for every $q_1 = \langle a_1, b_1 \rangle, q_2 = \langle a_2, b_2 \rangle \in W$ such that $p \in C_{q_1} \cap C_{q_2}$, we have: $q_1 \preceq_p^+ q_2$ if and only if $a_1 \leq a_2$. In particular, the $\preceq_p^+$ -ordering depends only on q_1, q_2 and not on p .

We claim that these U and W are the desired neighbourhoods. Indeed, if C_{q_1} and C_{q_2} intersect more than once in U , for $q_1, q_2 \in W$, then take two consecutive points of intersections p_1 and p_2 . Notice that if $q_1 \preceq_{p_1}^+ q_2$ then, because we assumed that C_{q_1} and C_{q_2} do not touch each other, we necessarily have $q_2 \preceq_{p_2}^+ q_1$. But this contradicts the fact that the $\preceq_p^+$ -ordering depends only on q_1 and q_2 (and not on p). $\square$

The following is a variation of a similar definition from [20].

Definition 1.7. Let $\mathcal{F} = \{C_q : q \in Q\}$ be a definable normal family of plane curves, all contained in $U \subseteq M^2$, where Q is an open subset of M^2 . We say that $\mathcal{F}$ is a *nice family* if the following hold:


- (i) For every $q_1, q_2 \in Q$, C_{q_1} and C_{q_2} intersect at most once in U .
- (ii) For every $q \in Q$, the curve C_q is the graph of a partial function $f_q : M \rightarrow M$.
- (iii) The partial function $F(a, b, x)$ which sends $\langle a, b, x \rangle \in Q \times M$ to $f_{\langle a, b \rangle}(x)$ is continuous in all variables and strictly monotone in each of its variables. Moreover,

in each of the variables, F has the same monotonicity behaviour, as the other two variables vary. E.g., if $F(a_1, b_1, -)$ is strictly increasing in the last variable, for some $\langle a_1, b_1 \rangle \in W$ then for all $\langle a, b \rangle \in W$, the function $F(a, b, -)$ is strictly increasing.

Corollary 1.8. *Let $\mathcal{F} = \{C_q : q \in Q\}$ be a 0-definable normal family of plane curves, where Q is a two-dimensional subset of M^2 . Assume that $p_0 \in C_{q_0}$, with $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$. Then there is an open neighbourhood $U \times W$ of $\langle p_0, q_0 \rangle$ such that the family $\mathcal{F}_{U,W} = \{C_q \cap U : q \in W\}$ is nice.*

Proof. Clause (i) of the definition of a nice family follows from Lemma 1.6, while the other clauses follow from genericity. $\square$

Definition 1.9. *Given a nice family of plane curves $\mathcal{F} = \{C_q : q \in Q\}$ and a definable curve $X \subseteq M^2$ we say that X is $\mathcal{F}$ -bounded at p if there are $q_1, q_2 \in Q$ such that $p \in C_{q_1}, C_{q_2}$ and $C_{q_1} \preceq_p X \preceq_p C_{q_2}$ (see figure)*



C is bounded at p by X and Y

The following is just Theorem 10.5 from [20]. The proof there uses no more than our definition of a nice family.

Fact 1.10. *Let $\mathcal{F}$ be a nice family of curves in $U \subseteq M^2$ and let $X \subseteq U$ be a definable curve that is $\mathcal{F}$ -bounded at every point. If p is generic in X then there exists a unique curve from $\mathcal{F}$ which touches X at p .*

In particular, if $X = C_q$ then the only curve from $\mathcal{F}$ that touches C_q at p is C_q itself.

Corollary 1.11. *Let $\mathcal{F}$ be a two-dimensional normal family of plane curves. Let $X \subseteq M^2$ be an A -definable curve, p generic in X (over A) and also generic in M^2 (over $\emptyset$). Then there are at most finitely many curves from $\mathcal{F}$ which touch X at p .*

Proof. If there were infinitely many touching curves in $\mathcal{F}$ we could find $q \in Q$ such that $\dim(p, q/A) = 3$ and C_q touches X at p . By 1.5, q is generic in Q over A . By Lemma 1.8, there is a neighbourhood $U \times W$ of $\langle p, q \rangle$ such that $\mathcal{F}_{U,W} = \{C_q : \langle p, q \rangle \in U \times W : p \in C_q\}$ is nice. Furthermore, we can choose U to be definable over independent parameters. It follows from 1.10 that $q \in \text{dcl}_{\mathcal{M}}(Ap)$, contradiction. $\square$

Definition 1.12. *Let $\mathcal{F}$ be a normal family of curves and X a definable curve in M^2 . For p a C^0 -point of X , let us denote by $\tau(X, p)$ the set of all $q \in Q$ such that C_q touches X at p .*

We let

$$\tau(X) = \bigcup \{\tau(X, p) : p \text{ a } C^0\text{-point of } X\} \subseteq Q.$$

We say that X is $\mathcal{F}$ -linear near p if there exist a neighbourhood U of p and $q \in Q$ such that for every $p' \in U \cap X$, we have $\tau(X, p') = \{q\}$.

If $\mathcal{F}$ is a nice family then X is $\mathcal{F}$ -linear near a generic p if and only if X equals near p to some C_q in $\mathcal{F}$. If $\mathcal{F}$ is not assumed to be nice then we just have the “only if” direction. Namely, it is possible that $C_q \in \mathcal{F}$ has curves from $\mathcal{F}$ (other than itself) touching it at p , in which case $X = C_q$ will not be $\mathcal{F}$ -linear.

Assume that $\mathcal{F}$ is a nice family of curves, X is $\mathcal{F}$ -bounded and not $\mathcal{F}$ -linear near any of its points. Then for any $p \in X$ for which $\tau(X, p) \neq \emptyset$, $\tau(X, p)$ and p are inter-definable in $\mathcal{M}$ over A . In particular, $\tau(X)$ is a definable 1-dimensional set. By corollary 1.11 this would still be true for normal $\mathcal{F}$ provided that there is no $p \in X$ such that $\{q \in Q : p \in C_q\}$ is two-dimensional. Furthermore, by 1.11, if $p \in X$ is generic and $q \in \tau(X, p)$ then q is generic in $\tau(X)$ over A .

Infinitesimals It is useful for much of what is coming next to introduce the notion of infinitesimals: Given $p \in M^n$, we denote by ν_p the *infinitesimal neighbourhood* of p , defined as follows: Given $\mathcal{M}^*$ an $|M|^+$ -saturated elementary extension of $\mathcal{M}$, ν_p is the intersection of all $\mathcal{M}$ -definable open neighbourhoods of p in $\mathcal{M}^*$. Notice that for $a = \langle a_1, \dots, a_n \rangle \in M^n$, we have $\nu_a = \nu_{a_1} \times \dots \times \nu_{a_n}$.

Even though the set ν_p is not definable, most of the statements involving ν_p appearing in this paper can be restated in a first order manner. For example, the statement “For every $q \in \nu_{q_0}$ the set $X \cap C_q \cap \nu_{p_0}$ is nonempty” is equivalent to “There exist neighbourhoods U of p_0 and W of q_0 such that for all $q \in W$, the set $X \cap C_q \cap U$ is nonempty”.

Lemma 1.13. *Let $\mathcal{F}$ be a nice family of plane curves in $U \subseteq M^2$, $X \subseteq U$ an A -definable curve that is $\mathcal{F}$ -bounded. Assume that $p_0 \in X$ is generic over A , that X is not $\mathcal{F}$ -linear near p_0 and that C_{q_0} touches X at p_0 . Let W_1, W_2 be the locally definably connected components of $M^2 \setminus \tau(X)$ at q_0 .*

Then, there is $i \in \{1, 2\}$, such that the following are true:

- (1) *Given any $q \in W_i \cap \nu_{q_0}$, the set $C_q \cap X \cap \nu_{p_0}$ is empty.*
- (2) *Given any $q \in W_{3-i} \cap \nu_{q_0}$, the set $C_q \cap X \cap \nu_{p_0}$ contains at least two points.*

Proof. Write $p_0 = \langle x_0, y_0 \rangle$, $q_0 = \langle a_0, b_0 \rangle$. As discussed above, $q_0 \in \tau(X)$ is generic over A so in particular it is a C^0 point of $\tau(X)$. Consider the continuous function $F(a, b, x) = f_{\langle a, b \rangle}(x)$, as given by the definition of nice families. Without loss of generality the domain of F is a product of intervals $I_1 \times I_2 \times I_3$ and its range is contained in the interval I_4 . Since p_0 is generic in X , the set X itself is also locally, near p_0 , the graph of a function, and without loss of generality we assume it is a function of x . Call this function $g(x)$. (If X is not a function of x , then, by genericity it is locally of the form $x = c$; in this case, notice that since $\mathcal{F}$ is nice, we can interchange the roles of x and y in $\mathcal{F}$ and consider it as a family of functions of y).

There are several cases to consider, but we will handle only one of them (the rest can be handled in a similar way). We assume that C_{q_0} touches X from above at p_0 . Namely, for all x near x_0 , we have $f_{q_0}(x) \geq g(x)$. Because p_0 is generic in X and C_{q_0} is the unique touching curve from $\mathcal{F}$ at p_0 , there is a neighbourhood U of p_0 such that for all $p \in U \cap X$, there is a touching curve $C_q \in \mathcal{F}$ to X at p and C_q touches X from above there.

Since q_0 is generic in $\tau(X)$ over A , we may assume that $\tau(X)$ is the graph of a function in the first variable.

Let us assume (by the definition of nice families) that $F(a, -, x)$ is strictly increasing in the second variable, near b_0 , for all a, x close to a_0, x_0 , respectively. Fix an interval I containing ν_{x_0} such that for all $\langle a, b \rangle \in \nu_{q_0}$ we have $I \subseteq \text{dom}(f_{a,b})$. It follows that for all $b > b_0$ and $x \in I$, we have $F(a_0, b, x) > F(a_0, b_0, x) > g(x)$. In particular, the curve $C_{a_0,b}$ lies above the curve X in some neighbourhood of p_0 and therefore the two curves do not intersect near p_0 (see Figure 3, with C_{q_0} for C_{a_0b}).




Figure 3:
The P universe




Figure 4:
The Q universe

Now, as we pointed out above, for all $q = \langle a, b \rangle \in \tau(X) \cap \nu_{q_0}$, the curve C_q touches X from above at some $p \in \nu_{p_0}$. If $q' = \langle a, b_1 \rangle \in \nu_{q_0}$ and $b_1 > b$ then by the same argument $C_{q'} \cap X \cap \nu_{p_0} = \emptyset$.

So, in the case we just considered, the local component W_1 of $M^2 \setminus \tau(X)$ which lies above the curve $\tau(X)$ satisfies: For all $q_1 \in W_1 \cap \nu_{q_0}$, $C_{q_1} \cap X \cap \nu_{p_0} = \emptyset$ (see Figures 3,4, with C_{q_1} for $C_{a_0b_1}$).

Let us consider the local component W_2 which lies below $\tau(X)$. If we take $b_1 < b_0$, and $\langle a_0, b_1 \rangle \in \nu_{q_0}$ then

$$(1) \quad f_{a_0,b_1}(x_0) < f_{a_0,b_0}(x_0) = g(x_0).$$

At the same time, by the previous argument there are $x_1 < x_0 < x_2$ both in ν_{x_0} such that $f_{a_0,b_0}(x_1) > g(x_1)$ and $f_{a_0,b_0}(x_2) > g(x_2)$. Therefore, by continuity considerations, if b_1 is sufficiently close to b_0 then there are $x'_1, x'_2 \in \nu_{x_0}$, $x'_1 < x_0 < x'_2$ such that

$$(2) \quad f_{a_0,b_1}(x'_1) > g(x'_1) \quad \text{and} \quad f_{a_0,b_1}(x'_2) > g(x'_2).$$

It follows by continuity from (1) and (2) that for some $x''_1, x''_2 \in \nu_{x_0}$ we have

$$(3) \quad f_{a_0,b_1}(x''_1) = g(x''_1) \quad \text{and} \quad f_{a_0,b_1}(x''_2) = g(x''_2).$$

Hence, X and C_{a_0,b_1} intersect at least twice in ν_{p_0} , for $b_1 < b$ and close to b_0 (see, again, Figures 3 and 4).

Repeating the same argument for every $q = \langle a, b \rangle \in \tau(X) \cap \nu_{q_0}$ and $\langle a, b_1 \rangle \in \nu_{q_0}$ with $b_1 < b$, we conclude that $|C_{a,b_1} \cap X \cap \nu_{p_0}| > 1$. $\square$

Lemma 1.14. *Let $\mathcal{F}$ be a nice family. Let $X \subseteq M^2$ be an A -definable curve, $p_0 \in X$ generic over A . If there is a curve from $\mathcal{F}$ that touches X at p_0 then X is $\mathcal{F}$ -bounded near p_0 .*

Proof. Assume that C_{q_0} touches X at p_0 . By Lemma 1.13 and its proof, there are q' arbitrarily close to q_0 and $p_1, p_2 \in \nu_{p_0}$ such that $C_{q'} \preccurlyeq_{p_1} X$ and $X \preccurlyeq_{p_2} C_{q'}$. But then the two sets

$$\{p' \in X : \exists q' C_{q'} \preccurlyeq_{p'} X\}$$

and

$$\{p' \in X : \exists q' X \preccurlyeq_{p'} C_{q'}\}$$

intersect ν_{p_0} nontrivially. Since these sets are definable just over the parameters defining X and $\mathcal{F}$, and since p_0 is generic in X over these parameters, the sets must contain all of $X \cap \nu_{p_0}$, hence X is $\mathcal{F}$ -bounded in a neighbourhood of p_0 . $\square$

Lemma 1.15. *Let $\mathcal{F}$ be a nice family of curves, X an A -definable curve, $p \in X$ generic over A and $q_0 \in Q$. If $p \in C_{q_0} \cap X$ then one and only one of the following holds:*

- (i) C_{q_0} touches X at p , or
- (ii) For every $q \in \nu_{q_0}$, we have $|C_q \cap X| = 1$.

Proof. Notice that by 1.13, clauses (i) and (ii) cannot hold simultaneously. Assume that C_{q_0} does not touch X at p .

Then C_{q_0} intersects both local components of $M^2 \setminus X$ in ν_p . By continuity, $C_{q'}$ intersects both components of $M^2 \setminus X$ in ν_p for all $q' \in \nu_{q_0}$. By the definable connectedness of $C_{q'}$ (or equivalently, the continuity of the function $f_{q'}$), every such curve $C_{q'}$ must intersect X in ν_p at least once.

Assume now, towards contradiction, that there exists $q \in \nu_{q_0}$ such that $C_q \cap X \cap \nu_p$ contains more than one point. We claim that X is $\mathcal{F}$ -bounded near p . There are two possibilities to consider:

(i) C_q touches X at one or more of these points of intersection. In this case, by Lemma 1.14 above, X is $\mathcal{F}$ -bounded in some open subset of ν_p . Because of the genericity of p (over the parameters defining $\mathcal{F}, X$), the curve X is $\mathcal{F}$ -bounded in a neighbourhood of p .

(ii) C_q does not touch X at any of their points of intersection. In this case, as we observed before, we can find $p_1, p_2 \in \nu_p$ such that $C_q \preccurlyeq_{p_1} X$ and $X \preccurlyeq_{p_2} C_q$. This was already shown to imply that X is $\mathcal{F}$ -bounded in ν_p .

Since X is $\mathcal{F}$ -bounded, by 1.10 there is a unique $q' \in Q$ such that $C_{q'}$ touches X at p . However, if we now replace Q by a smaller open neighbourhood of q_0 then the above argument shows that X is still $\mathcal{F}$ -bounded near p with respect to the restricted $\mathcal{F}$ and therefore there is still a curve in this family that touches X at p . This curve must therefore be C_{q_0} , contradicting the assumption that C_{q_0} does not touch X at p . $\square$

The last lemma can be seen as saying that in a nice family of curves, every curve that intersects X at a generic point either touches X or intersects it transversally.

Lemma 1.16. *Let $\mathcal{F}_0 = \{C_q : q \in Q_0\}$, $\mathcal{F}_1 = \{D_q : q \in Q_1\}$ be two 0-definable two-dimensional nice families of plane curves. Let q_0, q_1 be generic in Q_0, Q_1 , respectively, and $p_0 \in C_{q_0} \cap D_{q_1}$ such that $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = \dim_{\mathcal{M}}(p_0, q_1/\emptyset) = 3$. Assume that there exists some A -definable curve X such that p_0 is also generic in X over A . If C_{q_0} and D_{q_1} both touch X at p_0 then C_{q_0} and D_{q_1} touch each other at p_0 , and q_0 and q_1 are inter-definable over p_0 .*

Proof. Notice that if C_{q_0} and D_{q_1} touch X on opposite sides, then they clearly touch each other as well (see Figure 5). Assume then that the two curves touch X "from above". Because of the genericity of p_0 in X , for every $p \in \nu_{p_0}$, there is $q \in \nu_{q_0}$ such that C_q touches X from above at p . Similarly to the proof of 1.14, it follows that D_{q_1} is $\mathcal{F}_0$ -bounded near p_0 (see Figure 6). Moreover, even if we shrink Q_0 to a smaller neighbourhood of q_0 , the curve D_{q_1} is still $\mathcal{F}_0$ -bounded with respect to

this new family. As in the proof of Lemma 1.15, it follows (using 1.10), that C_{q_0} touches D_{q_1} at p_0 . (D_{q_1} plays the role of X from 1.10). By 1.11 (applied to the




Figure 5:
 C_{q_0}, D_{q_1} on opposite sides




Figure 6:
 D_{q_1} is $\mathcal{F}_0$ -bounded

family $\mathcal{F}_0$ and the curve D_{q_1}), if the two curves touch each other at p_0 then we have $q_0 \in \text{dcl}_{\mathcal{M}}(q_1, p_0)$. Similarly, $q_1 \in \text{dcl}(q_0, p_0)$. $\square$

1.4. Normal families and differentiability. *Throughout this subsection we assume that $\mathcal{M}$ expands a real closed field.*

Lemma 1.17. *Let $\{X_t : t \in T\}$ be a 0-definable normal family of plane curves, of dimension at least 2. Let $t_0 \in T$ be generic, $X = X_{t_0}$, and $p_0 = \langle x_0, y_0 \rangle \in X$ generic over t_0 . Then X is locally the graph of strictly monotone C^1 -function, $y = f_{t_0}(x)$. If d is the derivative of f_{t_0} at x_0 then $\dim_{\mathcal{M}}(d/p_0) = 1$.*

Proof. The first part follows from genericity together with the fact that $\dim_{\mathcal{M}} T \geq 2$ (otherwise, X could be of the form $x = c$). As for the second part, if $d \in \text{dcl}_{\mathcal{M}}(p_0)$ then there is a definable function $d(p)$, defined in a neighbourhood U_0 of p_0 , and there exists a neighbourhood T_0 of t_0 in T , such that for every $p \in U_0$ and every X_t containing p , with $t \in T_0$, the derivative of X_t at p equals to $d(p)$.

By the Uniqueness Theorem for differential equations in o-minimal structures (see Theorem 2.3 in [18]), for every X_t , $t \in T_0$, if $p_0 \in X_t$ then X_t and X_{t_0} coincide in some neighbourhood of p_0 . This contradicts normality. $\square$

Lemma 1.18. *Assume that $\mathcal{F} = \{C_q : q \in Q\}$ is a $\emptyset$ -definable two-dimensional normal family of plane curves. If $p_0 = \langle x_0, y_0 \rangle$ is generic in M^2 then for all but finitely many $q \in Q$, if $p_0 \in C_q$, then C_q is the graph of a C^1 -function $y = f_q(x)$ near p_0 . For every such q , $f'_q(x_0)$ and q are inter-definable over p_0 .*

Proof. For $p \in M^2$ let $\ell_p = \{q \in Q : p \in C_q\}$. The first part of the lemma follows from the fact that if q is generic in ℓ_{p_0} over p_0 then p_0 is generic in C_q over q (by 1.5).

Assume now $f_q(x)$ is C^1 near x_0 and that $d = f'_q(x_0)$. If there are infinitely many $q' \in \ell_{p_0}$ such that $f'_{q'}(x_0) = d$ then first of all there is $q_0 \in \ell_{p_0}$ such that $\dim_{\mathcal{M}}(q_0/p_0) = 1$ and $d = f'_{q_0}(x_0)$. Furthermore, because $\dim_{\mathcal{M}}(\ell_{p_0}) = 1$, we have $d \in \text{dcl}_{\mathcal{M}}(p_0)$. This contradicts Lemma 1.17 with C_{q_0} here taken as X . $\square$

Definition 1.19. *Let X be a definable curve. We say that X has rank k over A if there is a an A -definable normal family of curves $\{X_t : t \in T\}$ and $t_0 \in T$ generic over A , such that $X = X_{t_0}$ and $\dim_{\mathcal{M}}(T) = k$. We say that X has rank k if $A = \emptyset$.*

The following is easy to verify:

Remark 1.20. The notion of rank is well-defined, even locally. Namely, let $X \subseteq M^k$ be a definable curve of rank k over A , and U an A -definable open set with $U \cap X \neq \emptyset$. Then for every A -definable normal family of curves $\{X_t : t \in T'\}$, if there exists t in T' such that $X \cap U = X_t \cap U$ then $\dim_{\mathcal{M}}(T') \geq k$. In particular, $X \cap U$ has also rank k over A .

We can now establish the connection between the differential notion of tangency and that of touching curves.

Theorem 1.21. *Let $\mathcal{F} = \{C_q : q \in Q\}$ be an $\emptyset$ -definable normal family of plane curves of dimension greater than 1. Let X be an A -definable plane curve of rank greater than 1, $p_0 = \langle x_0, y_0 \rangle \in X$ generic over A , and d the derivative of the function associated to X at x_0 .*

If $p_0 \in C_{q_0}$ for $q_0 \in Q$ and if $f'_{q_0}(x_0) = d$ (in the notation above) then C_{q_0} and X touch each other at p_0 and $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$.

Proof. We assume that all sets are 0-definable. By 1.5, p_0 is generic in M^2 . It follows from 1.17 that $\dim_{\mathcal{M}}(d/p_0) = 1$, and from 1.18 it follows that $\dim_{\mathcal{M}}(q_0/p_0) = 1$, hence $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$.

We may assume now that X is the graph of a C^1 -function $y = h(x)$ near x_0 . By Theorem 1.8, we may assume that $\mathcal{F}$ is a nice family.

Assume towards contradiction that C_{q_0} does not touch X at p_0 . Say, we have $X \prec_{p_0} C_{q_0}$. It follows from continuity of the family that for every $q \in \nu_{q_0}$, there is $x > x_0$, $x \in \nu_{x_0}$, such that $f_q(x) > h(x)$. By 1.17, we can find $q \in \ell_{p_0} \cap \nu_{p_0}$ such that $f'_q(x_0) < d$ and therefore $C_q \prec_{p_0} X$. These two last facts, together with the continuity of f_q , imply that there is $x > x_0$, $x \in \nu_{x_0}$, such that $f_q(x_0) = h(x_0)$. In particular, $|C_q \cap X \cap \nu_{p_0}| > 1$, contradicting 1.15. $\square$

1.5. The Duality Theorem. For $\mathcal{F} = \{C_q : q \in Q\}$ a nice family of curves in $P \subseteq M^2$, with $Q \subseteq M^2$, we let $\mathcal{L} = \{\ell_p : p \in P\}$ be the dual family, defined by $q \in \ell_p \Leftrightarrow p \in C_q$. It is not hard to see that the monotonicity and continuity assumptions in the definition of a nice family imply that for every $p \in P$, either ℓ_p is empty or $\dim_{\mathcal{M}}(\ell_p) = 1$. Furthermore, because

$$q_1, q_2 \in \ell_{p_1} \cap \ell_{p_2} \Leftrightarrow p_1, p_2 \in C_{q_1} \cap C_{q_2},$$

for $p_1 \neq p_2$ we have $|\ell_{p_1} \cap \ell_{p_2}| \leq 1$. Finally, notice that if a definable function $G(x, y, z)$ is monotone and continuous in each variable separately then it is continuous as a function of all variables. We thus have:

Claim 1.22. *If $\mathcal{F}$ is a nice family then the dual family $\mathcal{L}$ is also nice.*

We can now establish the following duality:

Theorem 1.23. *Assume that $\mathcal{F}$ is a nice family of curves. Let $X \subseteq M^2$ be an $\mathcal{M}$ -definable curve and let p_0 be a generic point of X such that X is not $\mathcal{F}$ -linear near p_0 . If C_{q_0} touches X at p_0 then ℓ_{p_0} touches $\tau(X)$ at q_0 .*

Before proving the theorem, here is an example:

Example: Let $\mathcal{F}$ be the family of all affine lines in $\mathbb{R}^2$, $C_{a,b} = \{y = ax + b\}$. It is easy to see that the dual family $\mathcal{L}$ is also the family of affine lines. Take $X = \{y = x^2\}$.

Then $\tau(X) = \{\langle u, v \rangle : v = -u^2/4\}$. Given $p = \langle x_0, x_0^2 \rangle \in X$, the $\mathcal{F}$ -tangent curve at p is C_q , where $q = \langle 2x_0, -x_0^2 \rangle$. We also have $\ell_p = \{\langle u, v \rangle : v = -x_0u + x_0^2\}$.

The slope of ℓ_p equals everywhere to $-x_0$, which is exactly the slope of $\tau(X)$ at $q = \langle 2x_0, -x_0^2 \rangle$. Since q clearly belongs to ℓ_p it follows that ℓ_p is tangent to $\tau(X)$ at q .

Proof of theorem. Since p_0 and q_0 are inter-definable (over the parameters defining X), the point q_0 is generic in $\tau(X)$ as well.

Claim The curve ℓ_{p_0} is not equal to the curve $\tau(X)$ in a neighbourhood of q_0 .

Indeed, notice that if ℓ_{p_0} equals to $\tau(X)$ in a neighbourhood of q_0 then the same is true for all p in a relatively open $X_1 \subseteq X$ containing p_0 . But then all of these ℓ_p intersect each other at infinitely many points, contradicting the normality of $\mathcal{L}$.

Let W_1, W_2 be the two local components of $M^2 \setminus \tau(X)$ at q_0 . Without loss of generality, by Lemma 1.13, for every $q \in W_1 \cap \nu_{q_0}$, we have $C_q \cap X \cap \nu_{p_0} = \emptyset$. However, for every $q \in \ell_{p_0}$ we clearly have $p_0 \in C_q \cap X \cap \nu_{p_0}$. Hence, near q_0 , the entire curve ℓ_{p_0} is contained in W_2 , so ℓ_{p_0} touches $\tau(X)$ at q_0 . $\square$

Notice that the Duality Theorem implies, in the notation of the theorem, that if X is $\mathcal{F}$ -bounded near p_0 then $\tau(X)$ is $\mathcal{L}$ -bounded near q_0 (see Lemma 1.14).

Remark After discovering the above duality we found out that it has a classical analogue in projective geometry. In that setting a duality exists between an algebraic variety and the variety of all its tangent hyper-planes. See, e.g., [2].

We need the following corollary;

Corollary 1.24. *Let $\mathcal{F} = \{C_q : q \in Q\}$ be a $\emptyset$ -definable two dimensional normal family of curves in P , where $\dim P = 2$. Assume that q_0 is generic in Q , and $X \subseteq M^2$ is an A -definable curve. Assume that p_0, p_1 are generic elements of X and that X is not $\mathcal{F}$ -linear near any of them. If $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$ and C_{q_0} touches X at both p_0 and p_1 then $p_1 \in \text{dcl}_{\mathcal{M}}(p_0, q_0)$ (although X is A -definable!).*

Proof. We may clearly assume that $p_1 \notin \text{dcl}_{\mathcal{M}}(q_0)$. By 1.8, we may shrink P and Q , and obtain two nice families of curves $\mathcal{F}_0 = \{C_q^0 : q \in Q_0\}$ and $\mathcal{F}_1 = \{C_q^1 : q \in Q_1\}$, such that $C_{q_0}^0$ and C_{q_0} agree in a neighbourhood of p_0 and similarly, $C_{q_1}^1$ and C_{q_1} agree in a neighbourhood of p_0 . By choosing $\mathcal{F}_0$ and $\mathcal{F}_1$ properly we may absorb all new parameters and still assume that they are $\emptyset$ -definable.

By 1.23 the curves ℓ_{p_0} and ℓ_{p_1} touch $\tau(X)$ at q_0 . We now apply 1.16 to the families $\mathcal{L}_0$ and $\mathcal{L}_1$ (the dual families of $\mathcal{F}_0$ and $\mathcal{F}_1$), the points p_0, p_1 and $\tau(X)$, and conclude that $p_0 \in \text{dcl}_{\mathcal{M}}(q_0, p_1)$. $\square$

Remarks

- (1) Assume that $\mathcal{F}$ is a $\emptyset$ -definable normal family of plane curves in P (but $\mathcal{F}$ need not be a nice family), such that its dual family $\mathcal{L}$ is also normal. Assume also that $X \subseteq P$ is a definable curve of rank $k > 1$, p generic in X , C_{q_0} touches X at p_0 and $\dim(p_0, q_0/\emptyset) = 3$. By working locally, we can apply Theorem 1.23 and conclude that ℓ_{p_0} touches $\tau(X)$ at q_0 . However, in this case it is possible that there are finitely many such C_q , all touching X at p_0 , and hence ℓ_{p_0} will touch $\tau(X)$ at each one of these points q .

- (2) In many of the results in this section we add the extra assumption that $\dim_{\mathcal{M}}(p, q/\emptyset) = 3$. This assumption is indeed necessary, as the following example shows:

Let $C = \{\langle x, y \rangle \in \mathbb{R}^2 : y = \sqrt{|x|}\}$ and for $\langle a, b \rangle \in \mathbb{R}^2$ let $C_{a,b} = C + \langle a, b \rangle$. The family $\{C_{a,b} : \langle a, b \rangle \in \mathbb{R}^2\}$ is normal. If $X \subseteq \mathbb{R}^2$ is any C^1 -curve then for any $p_0 = \langle a, b \rangle \in X$, the curve $C_{a,b}$ touches X at p_0 , and clearly $q_0 = \langle a, b \rangle \in \text{dcl}_{\mathcal{M}}(p_0)$. In this case, if $C_{a,b}$ touches X at another point p_1 then it will not be true in general that $p_1 \in \text{dcl}_{\mathcal{M}}(p_0, q_0) = \text{dcl}_{\mathcal{M}}(p_0)$.

Finally, we have:

Lemma 1.25. *Assume that $\mathcal{F}$ is a $\emptyset$ -definable nice family of plane curves, parameterised by Q . Let X be a nowhere $\mathcal{F}$ -linear t -definable plane curve of rank $k > 1$ over A . For every $p_0 \in X$ generic over A, t , if C_{q_0} touches X at p_0 , then there exists a neighbourhood W of q_0 such that $\tau(X) \cap W$ is a curve of rank k over A .*

Proof. Because q_0 is generic in $Y = \tau(X)$ and $\mathcal{L}$ is a nice family, there are neighbourhoods U of p_0 and W of q_0 such that $p \mapsto \tau(X, p)$ induces a bijection of $X \cap U$ and $\tau(X) \cap W$, and moreover, by the Duality Theorem $\tau(\tau(X \cap U)) = X \cap U$.

Recall that $X = X_t$ for the A -definable family $\{X_t : t \in T\}$, where $\dim_{\mathcal{M}}(t/A) = k$, and consider the family $\{\tau(X_t \cap U) : t \in T\}$. We may assume that for all $t \in T$ we have $\tau(\tau(X_t \cap U)) = X_t \cap U$.

We claim that this last family must be normal. Indeed, if not then there are $t_1, t_2 \in T$ and an open set $W_1 \subseteq W$ such that $\tau(X_{t_1} \cap U) \cap W_1 = \tau(X_{t_2} \cap U) \cap W_1$. If we now apply τ to those sets then, by the Duality Theorem we will conclude, for some $U_1 \subseteq U$, that $X_{t_1} \cap U_1 = X_{t_2} \cap U_1$, contradiction. $\square$

1.6. Limit sets and a theorem of v.d. Dries. Somewhat surprisingly, the notion of limit sets plays a crucial role in our subsequent counting of intersection points of curves. Several different such notions of were studied extensively by v.d. Dries in [4]. The definition below, which is suitable for an arbitrary o-minimal structure (not necessarily over the reals) resembles most that of the Hausdorff limit, but is not restricted to families of compact, or even bounded sets.

Definition 1.26. *Consider a definable set $S \subseteq M^k \times M^n$. For $x \in Z_1 = \pi_1(S)$, the projection of S on the first k -coordinates, let $S_x = \{y \in M^n : \langle x, y \rangle \in S\}$. Let $\mathcal{S} = \{S_z : z \in Z_1\}$ be the corresponding family of subsets of M^n . Given $a < b$ in $M \cup \{\pm\infty\}$, and given a definable map $\gamma : (a, b) \rightarrow Z_1$, we let $\mathcal{S}(\gamma)$ be the set of all $y \in M^n$ such that for every open neighbourhood V of y and for every $\epsilon \in (a, b)$, there is $t \in (a, \epsilon)$ such that $(S_{\gamma(t)} \cap V) \neq \emptyset$.*

$\mathcal{S}(\gamma)$ is called the limit of $\mathcal{S}$ along $\gamma(t)$, as t tends to a .

Notice that equivalent definitions to the above are given by: $y \in \mathcal{S}(\gamma)$ iff for every $t > a$ in ν_a , $S_{\gamma(t)} \cap \nu_y \neq \emptyset$ iff there exists $t > a$ in ν_a such that $S_{\gamma(t)} \cap \nu_y \neq \emptyset$.

Here are some easy facts about limit sets:

(1) If every S_x has dimension k then the dimension of every limit set as above is at most k .

(2) If $\lim_{t \rightarrow a} \gamma(t) = z_0$ is a generic point of Z_1 (over the parameters defining $\mathcal{S}$) then $\mathcal{S}(\gamma) = \text{cl}(S_{z_0}) = \text{cl}(S) \cap (\{z_0\} \times M^n)$. In particular, for any other definable $\gamma' : (a', b') \rightarrow Z_1$ such that $\lim_{t \rightarrow a'} \gamma'(t) = z_0$, we have $\mathcal{S}(\gamma) = \mathcal{S}(\gamma')$.

(3) Assume that $X \subseteq Z_1$ is a definable one-dimensional set, $p \in \text{cl}(X)$. Then there are finitely many ways to approach p in X (depending on the number of local

components of $X \setminus \{p\}$). We write $\mathcal{S}(X, p)$ for the union of the finitely many possible limits sets $\mathcal{S}(\gamma)$ as $\gamma(t)$ tends to p in X , and call this set *the limit set of $\mathcal{S}$ along X at p* . If p is a generic point in X then $\mathcal{S}(X, p)$ is just $\text{cl}(S_p)$.

The following theorem is due to van den Dries (the complete - unpublished - proof of which appears in [5], and is similar to the proof of the analogues theorem in §9 of [4]):

Theorem 1.27. *Assume that $\mathcal{M}$ expands a real closed field. Given a definable family $\mathcal{S}$ as above, the family*

$$\tilde{\mathcal{S}} = \{\mathcal{S}(\gamma) \mid \gamma : (a, b) \rightarrow Z_1 \text{ definable} \}$$

is definable in $\mathcal{M}$ and $\dim(\tilde{\mathcal{S}}) \leq \dim Z_1$. If furthermore all the sets in $\mathcal{S}$ are closed then $\dim(\tilde{\mathcal{S}} \setminus \mathcal{S}) < \dim Z_1$ (By $\dim(\tilde{\mathcal{S}})$ we mean the smallest possible dimension for a set of parameters for $\tilde{\mathcal{S}}$).

In this paper we will be using only the first part of the theorem. Since the proof does not appear elsewhere, we present in Appendix A, with the author's permission, his proof of the theorem.

1.7. Special points. *In this section we assume that $\mathcal{F} = \{C_q : q \in Q\}$ is a 0-definable two-dimensional normal family of curves in $P \subseteq M^k$, $\dim P = 2$ (but P is not necessarily contained in the plane), such that its dual family $\mathcal{L}$ is also normal.*

A few words of explanations are in place regarding these assumptions.

- (1) If $\mathcal{F}$ is an (almost) normal two-dimensional family of curves, its dual family need not consist only of curves, i.e. for $p \in P$ the set ℓ_p of $\mathcal{F}$ -curves through p may be 2-dimensional. It is easy to check that by normality the set $\hat{P} := \{p \in P : \dim \ell_p = 2\}$ is finite. Restricting to $P \setminus \hat{P}$, the dual to $\mathcal{F}$ is also a family of curves.
- (2) If $\mathcal{F}$ is a normal two-dimensional family of curves whose dual is a family of curves as well, it is easy to check that the dual is almost normal. However, it need not be normal. In the application in Section 2 the family $\mathcal{F}$ will be definable in a strongly minimal structure. We leave it as simple exercise to check that in that case replacing p with p/E for an appropriate definable equivalence relation E both $\mathcal{F}$ and its dual are normal. See also Fact 2.4.
- (3) There is no harm assuming that $\mathcal{F}$ is only almost normal (whose dual is a family of curves), in which case its dual is easily checked to be almost normal as well.

The next definition, which is quite technical, is motivated by the following argument: Our ultimate goal is to prove that the existence of an almost normal two-dimensional $\mathcal{N}$ -definable family $\mathcal{F}$ of plane curves contradicts the stability of $\mathcal{N}$. The idea is to start with a curve X , and a curve C_{q_0} from $\mathcal{F}$ touching X at a generic point, p_0 . By Lemma 1.13, there are curves arbitrarily close to C_{q_0} which either intersect X nowhere near p_0 or intersect it twice there. If there were no other intersection points of these curve with X , this gap in the number of intersection points with X can be seen to contradict stability. However, the problem is that the curves near C_{q_0} might intersect X elsewhere and in order to control the total number of intersection points we need to investigate what happens near those points. Such points will be called *special for (q_0, X)* .

Definition 1.28. Let $X \subseteq P$ be a definable curve. Given $q \in Q$, a point $p \in M^k$ is called special for (q, X) if there exist $q' \in \nu_q$, and $p' \in \nu_p$ such that $p' \in X \cap C_{q'}$ (equivalently, for every neighbourhoods U, V of q, p , respectively, there exist $q' \in U, p' \in V$, such that $p' \in X \cap C_{q'}$).

A similar definition can be given even if some of the coordinates of p are taken to be $\pm\infty$ (and then $\nu_{+\infty}, \nu_{-\infty}$ have an obvious meaning).

The point p in the above definition can be seen as an asymptotic direction of X , in the sense of the family $\mathcal{F}$, and the curve C_q can be seen as an “ $\mathcal{F}$ -asymptote of X at p ”. For example, if $\mathcal{F}$ is the family of affine lines in $\mathbb{R}^2$, given by $C_{a,b} = \{y = ax + b\}$, and X is the curve $y(x-1) = 1$ then the point $p = \langle 1, +\infty \rangle$ is special for $(\langle 0, 1 \rangle, X)$ (that is, the curve $y = 1$ is asymptotic to X at $(1, +\infty)$). Indeed, curves of the form $y = 1 + \epsilon$ will intersect X near $+\infty$.

The following are easy observations concerning special points:

- (1) If $p \in C_q \cap X$ then clearly p is a special point for (q, X) .
- (2) If $p \in M^k$ is a special point for (q, X) then $p \in \text{cl}(X) \subseteq \text{cl}(P)$. However, if $\mathcal{F}$ is not given continuously, it is possible that p is not in $\text{cl}(C_q)$.
- (3) The set of special points for (q, X) is clearly definable.
- (4) If p is a special point for (q, X) then, by curve selection, there exist an interval (a, b) and a definable curve $\gamma : (a, b) \rightarrow Q$ tending to q as t tends to a , such that p is in the limit set $\mathcal{F}(\gamma)$. In particular, if q is generic in Q then $p \in \mathcal{F}(\gamma) = \text{cl}(C_q)$ (see (2) of the previous section).

The following lemma established the connection between the notions of a special point and that of a limit set:

Lemma 1.29. Given $\mathcal{F}$ and a curve X , the point p is a special point for (q, X) if and only if q belongs to the limit set $\mathcal{L}(X, p)$, of the dual family $\mathcal{L}$ along X at p (see (3) of the previous subsection)

Proof. By definition, p is special for (q, X) if and only if $X \cap \nu_p \cap C_{q'} \neq \emptyset$ for some $q' \in \nu_q$, if and only if $\ell_{p'} \cap \nu_q \neq \emptyset$ for some $p' \in X \cap \nu_p$. Using the remark immediately after the definition of limit sets, this is equivalent to q belonging to the limit set of $\mathcal{L}$ at p , along the curve X . $\square$

Lemma 1.30. Let q_0 be generic in Q and let $X \subseteq P$ be a definable curve not $\mathcal{F}$ -linear near any of its points. Then:

- (1) There are at most finitely many points $p_1, \dots, p_r$ (including possibly points with coordinates in $\pm\infty$), which are special for (q_0, X) .
- (2) There are definable open neighbourhoods $U_1, \dots, U_r$ of $p_1, \dots, p_r$, respectively, and an open neighbourhood W of q_0 such that for every $q \in W$, every intersection point of C_q with X is contained in one of the U_i .

Proof. (1) As was remarked above, if p is special for (q_0, X) then there is a definable curve $\gamma : (a, b) \rightarrow Q$, tending to q_0 as t tends to a , such that $p \in \mathcal{F}(\gamma) = \text{cl}(C_{q_0})$. Now, if there were infinitely many special points for (q_0, X) then infinitely many of them were already in $C_{q_0} \cap X$ and in particular, C_{q_0} would equal X in some neighbourhood, contradicting our choice of X .

(2) If the statement failed then, by curve selection, we would get a definable curve γ tending to q_0 with points in $C_{\gamma(t)} \cap X$ tending away from $p_1, \dots, p_r$. By definition,

if p is a limit point for the set $C_{\gamma(t)} \cap X$ then it is a special point for (q_0, X) (some of the coordinates of p could be $\{\pm\infty\}$). This would contradict the assumption that $p_1, \dots, p_r$ were all the special points for (q_0, X) . $\square$

The following technical lemma will play a very important role in our subsequent proof of the main theorem.

Lemma 1.31. *Let $X \subseteq P$ be an A -definable curve of rank $k > 2$, p_0 generic in X and assume that for $q_0 \in Q$, the curve C_{q_0} touches X at p_0 (here we identify P locally with an open set in M^2), and in addition, $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$. If p is a special point for (q_0, X) then $\dim_{\mathcal{M}}(p/A) = 1$. In particular, p is a generic point of X .*

Proof. As was pointed out above, $p \in \text{cl}(X)$ and hence $\dim_{\mathcal{M}}(p/A) \leq 1$. By 1.6, we may assume that in a neighbourhood U of p_0 and W of q_0 , the family $\mathcal{F}$ is nice. Consider $\tau(X \cap U)$ and recall that q_0 is generic in $\tau(X \cap U)$ over A (by the Duality Theorem).

Assume, towards a contradiction, that $\dim_{\mathcal{M}}(p/A) = 0$. Hence, q_0 is still generic in $\tau(X \cap U)$ over Ap . As we pointed out in (3) above, q_0 is in the Ap -definable limit set $\mathcal{L}(X, p)$. But then there is a neighbourhood $W_1 \subseteq W$ of q_0 in $\tau(X \cap U)$, such that every q in $\tau(X \cap U) \cap W_1$ belongs to $\mathcal{L}(X, p)$. The set $\mathcal{L}(X, p)$ has dimension one and therefore $\tau(X \cap U)$ equals to $\mathcal{L}(X, p)$ in some open set $W_2 \subseteq W_1$.

The curve $\mathcal{L}(X, p)$ is a member of the family of limit sets $\tilde{\mathcal{L}}$, hence by Theorem 1.27, its dimension is at most 2. It follows that the rank of $\mathcal{L}(X, p) \cap W_2 = \tau(X \cap U) \cap W_2$ is at most 2. However, by 1.25 (after possibly shrinking W_2), the rank of $\tau(X \cap U) \cap W_2$ equals that of X , contradicting our assumption that $\text{Rank } X \geq 3$. $\square$

Remark The above lemma implies that, under the same assumptions, a point in $\text{Fr}(X)$ cannot be special for (q_0, X) (see Figure 7). Similarly, it can be shown that such special points cannot have $\pm\infty$ as one of their coordinates. This can be seen as saying that a generic tangent curve to X cannot be asymptotic to X at $\pm\infty$.




Figure 7:
A problematic special point

We will need the following lemma:

Lemma 1.32. *Let $\mathcal{F}$, $\mathcal{L}$, X , p_0 and q_0 be as in the last lemma. Assume that p is a special point for (q_0, X) such that $\dim_{\mathcal{M}}(p/p_0, q_0) = 1$. Then, for all $q \in \nu_{q_0}$, the curve C_q intersects X exactly once in ν_p .*

Proof. By Lemma 1.31, p is generic in X and by Lemma 1.24, C_{q_0} does not touch X at p .

The fact that p is a special point for (q_0, X) implies (see (4) after the definition of special points) that p is in $\text{cl}(C_{q_0})$. However, since we assume that $\dim_{\mathcal{M}}(p/q_0) = 1$, p must belong to C_{q_0} . We now apply Lemma 1.15 to $\mathcal{F}_{U,W}$ for some neighbourhood U of p and W of q_0 , such that $\mathcal{F}_{U,W}$ is nice. $\square$

2. STABLE ONE DIMENSIONAL THEORIES IN O-MINIMAL STRUCTURES

We now return to the original setting of the main theorem. Namely, $\mathcal{M}$ is an o-minimal structure (expanding a real closed field), $\mathcal{N}$ is definable in $\mathcal{M}$ such that $\dim_{\mathcal{M}}(N) = 1$. In this section we treat the case that $\mathcal{N}$ is stable. First note that by cell decomposition, and the fact that we assume that $\mathcal{M}$ expands a field, we may assume that N is a subset of M (later on, in Appendix B, we will show why this can be assumed even without the field assumption)

We will need the following facts about stable theories interpretable in o-minimal structures:

Fact 2.1. *Assume $\mathcal{N}$ is a stable structure interpretable in $\mathcal{M}$. Then:*

- (1) $\mathcal{N}$ is superstable and its U -rank is at most the o-minimal dimension of N (see [7]).
- (2) If $U(\mathcal{N}) = 1$ and $\mathcal{N}$ is non trivial then $R^\infty(\mathcal{N}) = 1$. If in addition $\mathcal{N}$ is non locally modular then $RM(\mathcal{N}) = 1$ (see Chapter 2, Corollary 3.3 in [22]).

Our goal in this section is to prove;

Theorem 2.2. *If $\mathcal{N}$ is stable then the acl-geometry is necessarily locally modular.*

Proof. By the properties above, $U(\mathcal{N}) = 1$. We assume towards contradiction that the acl-geometry is not locally modular. Hence, by the same properties, we may assume that $\mathcal{N}$ is strongly minimal. Our strategy is to find a strongly minimal field definable in $\mathcal{N}$. This leads to an immediate contradiction since on the one hand, by a classical result of MacIntyre's any such field must be algebraically closed (see, e.g. [23]) whereas any strongly minimal field definable in $\mathcal{N}$ is necessarily 1-dimensional in the o-minimal sense and hence it is real closed (see [3]).

Recall that we use $\dim_{\mathcal{N}}$ to denote the dimension of tuples and sets in the sense of the geometric structure $\mathcal{N}$. Since $\mathcal{N}$ is strongly minimal this is the same as Morley Rank.

Non-local modularity of $\mathcal{N}$ supplies us with a two-dimensional family of $\mathcal{N}$ -definable plane curves, to which we intend to apply the results of the previous section. So we remind:

Definition 2.3. A definable (possibly in $\mathcal{N}^{eq}$) *pseudoplane* is given by two sets P, Q , together with a definable set $F \subseteq P \times Q$, such that the following holds:

- (i) $MR(P) = MR(Q) = 2$.
- (ii) For every $q \in Q$, $MR(F(P, q)) = 1$ (where $F(P, q) = \{p \in P : \langle p, q \rangle \in F\}$).
- (iii) For every $p \in P$, $MR(F(p, Q)) = 1$ (where $F(p, Q) = \{q \in Q : \langle p, q \rangle \in F\}$).
- (iv) For every $q_1 \neq q_2$ in Q , $F(P, q_1) \cap F(P, q_2)$ is finite.
- (v) For every $p_1 \neq p_2$ in P , $F(p_1, Q) \cap F(p_2, Q)$ is finite.

Given a pseudoplane as above, $p \in P, q \in Q$, we let $C_q = F(P, q)$ and $\ell_p = F(p, Q)$. We also let $\mathcal{F}$ denote the family of curves $\{C_q : q \in Q\}$ in P , and let $\mathcal{L}$ denote the dual family.

The following can be found in [25] (or see Proposition 1.7 on p.155 of [22]):

Fact 2.4. *If $\mathcal{N}$ is any strongly minimal nonlocally modular structure, then a pseudoplane is interpretable in $\mathcal{N}$. Moreover, (see, e.g. [11]), P may be taken to be a subset of N^2 (but without EI, we cannot assume at the same time that Q is a subset of N^2 as well).*

From now on we work with a fixed such pseudoplane that we assume is interpretable in $\mathcal{N}$. From the point of view of $\mathcal{M}$, the sets P and Q are 2-dimensional definable (using $\mathcal{M}$ -definable choice for $\mathcal{N}$) sets, with $\mathcal{F}$ and its dual $\mathcal{L}$ definable as well. Clearly, both $\mathcal{F}$ and $\mathcal{L}$ are normal families. We may assume that P, Q and each generic curve in $\mathcal{F}$ and $\mathcal{L}$ are of Morley degree 1. We may also assume that P, Q and $\mathcal{F}$ are all $\emptyset$ -definable.

The main stage in defining a field in $\mathcal{N}$ is to define a group. This will be achieved in two steps.

Step I Constructing an $\mathcal{N}$ -definable curve $X \subseteq P$ of rank $k \geq 3$, which has a sufficiently generic curve from $\mathcal{F}$ touching it at some generic p_0 .

We fix $p_0 = \langle x_0, y_0 \rangle$ an $\mathcal{M}$ -generic point in P . and 3 points:

$$p_3 = \langle z_2, y_0 \rangle, p_2 = \langle z_2, z_1 \rangle p_1 = \langle x_0, z_1 \rangle,$$

such that x_0, y_0, z_1, z_2 are $\mathcal{M}$ -generic and $\mathcal{M}$ -independent in N , all close to x_0 (we will see later how close we need them further to be).

We let Q_i , $i = 0, 1, 2, 3$, be the set of all $q \in Q$ such that $p_i \in C_q$. So, each Q_i is $\mathcal{N}$ -definable and has dimension (and Morley rank) one. We may assume that each Q_i has Morley degree one. For $q_1 \in Q_1, q_2 \in Q_2$ and $q_3 \in Q_3$ we let

$$C_{q_3 q_2 q_1} = C_{q_3} \circ C_{q_2}^{-1} \circ C_{q_1}.$$

Notice that if q_1, q_2, q_3 is an $\mathcal{M}$ -generic and $\mathcal{M}$ -independent sequence with $q_i \in Q_i$, then p_0 is $\mathcal{M}$ -generic in $C_{q_3 q_2 q_1}$ (over $q_3 q_2 q_1$). Fix $X = C_{q_3 q_2 q_1}$ for such an $\mathcal{M}$ -generic, $\mathcal{M}$ -independent sequence of q_i .

Case 1 The rank of X over p_1, p_2, p_3 is smaller than 3.

In that case, we obtain an $\mathcal{N}$ -definable infinite field. To do this, recall the following theorem of Hrushovski's (see, e.g., [1]):

Fact 2.5. *Suppose that M is a strongly minimal structure and there exists is a collection of tuples such that $\text{MR}(g_1, g_2, g_3, b_1, b_2, b_3) = 5$ and:*

- $\text{MR}(g_i) = 2$ and $\text{MR}(b_i) = 1$ for all $1 \leq i \leq 3$.
- The g_i are pairwise independent but $g_3 \in \text{acl}(g_1, g_2)$.
- $\text{MR}(g_1, b_1, b_2) = \text{MR}(g_2, b_2, b_3) = \text{MR}(g_3, b_1, b_3) = 3$.

Then a strongly minimal field is interpretable in M . We call such a collection of tuples a field configuration.

We will show that under the above assumptions we can find a field configuration. Note, first, that the rank of X over p_1, p_2, p_3 is at least 2 (because fixing q_2, q_3 we obtain, moving q_1 , a family of the same dimension as Q). So assume that the rank of X is precisely 2. In particular this is the rank of $C_{q_2}^{-1} \circ C_{q_1}$, i.e. $\text{MR}(Cb(C_{q_2}^{-1} \circ C_{q_1})) = 2$. Let $x \in N$ be generic over q_1, q_2 and choose $q_1(x) \in C_{q_1}(x) \setminus \text{acl}_{\mathcal{N}}(x)$, where $C_{q_1}(x) := \{y : \langle x, y \rangle \in C_{q_1}\}$. Let $q_2^{-1} q_1(x)$ be any point in $(C_{q_2}^{-1} \circ C_{q_1})(x) \setminus \text{acl}_{\mathcal{N}}(x)$. Then

$$Q := (Cb(C_{q_1}), Cb(C_{q_2}), Cb(C_{q_2} \circ C_{q_1}^{-1}), x, q_1(x), q_2^{-1} q_1(x))$$

form a field configuration over p_1, p_2, p_3 . To see this, note first that $\text{MR}(Q) = 5 = \text{MR}(q_1, q_2, x)$ since all the other element in Q are, by construction, algebraic over them. The first two bullets in the definition of a field configuration being immediate from the construction of Q we only have to check the third condition. But this

follows from the fact that x is generic and (e.g.) C_{q_1} is a generic curve through $(x, q_1(x))$ whence $\text{MR}(Cb(C_{q_1})/(x, q_1(x))) = 1$. The rest follows in a similar way. As we have already said, the existence of a strongly minimal (hence $\mathcal{M}$ -1-dimensional) field interpretable in $\mathcal{N}$ leads to a contradiction with the stability of CN .

So we are reduced to:

Case 2 The rank of X over p_1, p_2, p_3 is at least 3.

Let R denote the field underlying the structure $\mathcal{M}$. Because p_0, p_1, p_2, p_3 are generic in $X, C_{q_1}, C_{q_2}, C_{q_3}$, respectively, the curves are all C^1 -functions in neighbourhoods of those points. If we take x_0, y_0, z_1, z_2 sufficiently close to each other and q_1, q_2, q_3 sufficiently close to each other then by continuity arguments, the derivative of X at p_0 is close to those of the C_{q_i} at the p_i . Therefore, by Lemma 1.18 and continuity, we can find $q_0 \in Q_0$ such that C_{q_0} has the same derivative as X at the point p_0 . By Theorem 1.21, the curve C_{q_0} touches X at p_0 , and $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$.

End of Step I

Step II Counting Intersection points of X with $\mathcal{F}$ -curves.

We now fix the curve X , p_0 generic in X and C_{q_0} touching X at p_0 as given by Step I. We will investigate the intersection with X of $\mathcal{F}$ -curves close (in Q) to q_0 . By Lemma 1.30, all intersection points of such curves with X appear near the special points for (q_0, X) . By the first part of the same lemma, there are at most finitely many such special points $s_0 = p_0, s_1, \dots, s_r$

The General Case: All s_i , except s_0 , have $\dim_{\mathcal{M}}(s_i/p_0, q_0) = 1$.

By Lemma 1.30 there exist neighbourhoods W of q_0 and pairwise-disjoint U_i of s_i , $i = 0, \dots, r$, respectively, such that for every $q \in W$, every intersection point of C_q with X is in one of the U_i . Furthermore, by Lemma 1.32 we can choose W and the U_i in such a way that for each $q \in W$ and $i \neq 0$, the curve C_q intersects X in U_i exactly once.

Now, by Lemma 1.13, we can find $q_1, q_2 \in W$ both $\mathcal{N}$ -generic over all the data, such that, after possibly shrinking U_0 further, we have $C_{q_1} \cap X \cap U_0 = \emptyset$ while $|C_{q_2} \cap X \cap U_0| > 1$.

It follows that

$$|X \cap C_{q_1}| = r,$$

while

$$|X \cap C_{q_2}| \geq r + 2.$$

This contradicts the assumption that the Morley degree of Q is 1.

The Special Case: There exists s_i , $i = 1, \dots, r$ such that $\dim_{\mathcal{M}}(s_i/p_0, q_0) = 0$. The crucial part of the proof is:

Lemma 2.6. *If, under the above assumptions, $\dim_{\mathcal{M}}(s_1/p_0, q_0) = 0$ then a group is interpretable in $\mathcal{N}$.*

Proof. We still work with the 4 points

$$p_0 = \langle x_0, y_0 \rangle, p_1 = \langle x_0, z_1 \rangle, p_2 = \langle z_2, z_1 \rangle, p_3 = \langle z_2, y_0 \rangle,$$

such that x_0, y_0, z_1, z_2 are $\mathcal{M}$ -generic and $\mathcal{M}$ -independent. As above, we let Q_i , $i = 1, 2, 3$, be the set of all q such that $p_i \in C_q$. So, each Q_i is $\mathcal{N}$ -definable and has dimension one. We denote elements of Q_2 by r, r_1, r_2 etc.

For $q_1 \in Q_1, r \in Q_2$ and $q_3 \in Q_3$ we use $C_{q_3 r q_1} = C_{q_3} \circ C_r^{-1} \circ C_{q_1}$ as before. We denote the composition curves $C_{q_3} \circ C_{r_2}^{-1}$ and $C_{r_2}^{-1} \circ C_{q_1}$ by $C_{q_3 r_2}$ and $C_{r_2 q_1}$, respectively.

We fix $\mathcal{M}$ -independent $\hat{q}_3, \hat{r}, \hat{q}_1$ $\mathcal{M}$ -generic in Q_1, Q_2, Q_3 , respectively, and, as before, $X = C_{\hat{q}_3 \hat{r} \hat{q}_1}$, and C_{q_0} touches X at p_0 , with $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$. We assume also that there is $s_1 \in P$ special for (q_0, X) such that $s_1 \in \text{dcl}_{\mathcal{M}}(p_0, q_0)$. By Lemma 1.31, s_1 is generic in X over the parameters defining X (so $s_1 \notin \text{Fr}(X)$).

Our goal now is to construct a group configuration in $\mathcal{N}$. First, since

$$\dim_{\mathcal{M}}(\hat{q}_1, \hat{r}, \hat{q}_3/p_0, p_1, p_2, p_3) = 3,$$

there are relatively open neighbourhoods $Q'_1 \subseteq Q_1$, $Q'_2 \subseteq Q_2$ and $Q'_3 \subseteq Q_3$ of $\hat{q}_1, \hat{r}, \hat{q}_3$ respectively (definable in $\mathcal{M}$!!) such that:

$$(*) \quad \begin{aligned} &\text{for every } q_1 \in Q'_1, r \in Q'_2, q_3 \in Q'_3, \\ &\text{if } C_{q_0} \text{ touches } C_{q_3 r q_1} \text{ at } p_0 \text{ then } s_1 \in C_{q_3 r q_1} \end{aligned}$$

This can be done because touching is an $\mathcal{M}$ -definable notion and there is a p_0 -definable function sending q_0 to s_1 .

We first claim that for $r_1, r_2 \in Q_2$ which are $\mathcal{M}$ -independent and $\mathcal{M}$ -generic (over all mentioned parameters), the curve $C_{r_1 r_2} := C_{r_1} C_{r_2}^{-1}$ (we will use this notation whenever $r_1, r_2 \in Q_2$) is C^1 in a neighbourhood of $\langle z_1, z_1 \rangle$: Indeed, because $\langle z_2, z_1 \rangle$ is generic in both C_{r_1}, C_{r_2} , it is C^1 in both. It is therefore sufficient to see that the germ of $C_{r_1 r_2}$ at $\langle z_1, z_1 \rangle$ equals the composition of the germ of C_{r_1} at $\langle z_2, z_1 \rangle$ with the germ of $C_{r_2}^{-1}$ at $\langle z_1, z_2 \rangle$. For that, we need to verify that there is no $z'_2 \neq z_2$ such that $\langle z'_2, z_1 \rangle \in \text{cl}(C_{r_1}) \cap \text{cl}(C_{r_2})$.

If such $\langle z'_2, z_1 \rangle$ were in $\text{Fr}(C_{r_i})$ then z_1 would not be generic over the parameters defining C_{r_1} , contradiction. So we may assume that $\langle z'_2, z_1 \rangle \in C_{r_1} \cap C_{r_2}$, with $z'_2 \neq z_2$. Since the set $\{z : \langle z, z_1 \rangle \in C_{r_1}\}$ is finite we have $z_2, z'_2 \in \text{dcl}_{\mathcal{M}}(z_1, r_1)$. Because r_2 was chosen to be generic in Q_2 over z_1, r_1, p_2 , we get infinitely many $r \in Q_2$ such that $\langle z_2, z_1 \rangle, \langle z'_2, z_1 \rangle \in C_r$. This contradicts the normality of $\mathcal{L}$.

We can now describe the idea for the rest of the proof.

The idea: Generically, we will be able to recognise in $\mathcal{N}$ when $C_{r_1 r_2}$ and $C_{r_3 r_4}$ have the same derivative at $\langle z_1, z_1 \rangle$, for $r_1, r_2, r_3, r_4 \in Q_2$. This is done via the following relation:

$C_{r_1 r_2}$ and $C_{r_3 r_4}$ are tangent to each other at $\langle z_1, z_1 \rangle$ iff there are $q_1, q'_1 \in Q'_1$ and $q_3, q'_3 \in Q'_3$ such that the curves $C_{q_3 r_1 q_1}$, $C_{q_3 r_2 q'_1}$, $C_{q'_3 r_3 q_1}$ and $C_{q'_3 r_4 q'_1}$ all touch C_{q_0} at p_0 .

Indeed, notice that if $C_{q_3 r_1 q_1}$ and $C_{q'_3 r_3 q_1}$ touch C_{q_0} at $p_0 = \langle x_0, y_0 \rangle$ then in particular they have the same derivative at p_0 and therefore $C_{q_3 r_1}$ and $C_{q'_3 r_3}$ also have the same derivative at $\langle z_1, y_0 \rangle$. Similarly, $C_{q_3 r_2}$ and $C_{q'_3 r_4}$ have the same derivative at $\langle z_1, y_0 \rangle$. By composition, it now follows that $C_{r_1 r_2}$ and $C_{r_3 r_4}$ have the same derivative at $\langle z_1, z_1 \rangle$. Why is the above relation $\mathcal{N}$ -definable? By (*), for independent

generic q_1, r, q_3 in Q'_1, Q'_2, Q'_3 , respectively, tangency of $C_{q_3 r q_1}$ to C_{q_0} at p_0 can be detected by the fact that s_1 belongs to $C_{q_3 r q_1}$.

Finally, using the fact that there is an underlying real closed field the above tangency relation will allow us to obtain a group configuration in $\mathcal{N}$.

The actual argument : For $r_1, r_2, r_3, r_4 \in Q_2$ we define the relation

$$\begin{aligned} T(r_1, r_2, r_3, r_4) \Leftrightarrow \\ \text{there are } q_1, q'_1 \in Q_1 \text{ and } q_3, q'_3 \in Q_3 \text{ such that} \\ s_1 \in C_{q_3 r_1 q_1} \cap C_{q_3 r_2 q'_1} \cap C_{q'_3 r_3 q_1} \cap C_{q'_3 r_4 q'_1} \end{aligned}$$

T is clearly definable in $\mathcal{N}$. The main properties of T are:

Claim 2.7. Denote $A := \{s_1, p_0, p_1, p_2, p_3, q_0\}$ then:

- (1) For $\langle r_1, r_2 \rangle$ and $\langle r_3, r_4 \rangle$, each $\mathcal{M}$ -generic in $Q'_2 \times Q'_2$ and sufficiently close to $\hat{r}$, if the curves $C_{r_1 r_2}$ and $C_{r_3 r_4}$ have the same derivative at $\langle z_1, z_1 \rangle$ then $\langle r_1, r_2, r_3, r_4 \rangle \in T$.
- (2) If $\langle r_1, r_2, r_3 \rangle$ is an $\mathcal{N}$ -generic triple of elements from Q_2 then there exists $r_4 \in Q_2$, $r_4 \in \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 A)$, such that $\langle r_1, r_2, r_3, r_4 \rangle \in T$.

The proof of (1): Assume that $C_{r_1 r_2}$ and $C_{r_3 r_4}$ have the same derivative at $\langle z_1, z_1 \rangle$ (and the r_i are close to each other).

Pick $q_3 \in Q_3$ to be $\mathcal{M}$ -generic over r_1, r_2, r_3, r_4, q_0 and close to $\hat{q}_3$. Because r_1, r_3 are close to each other, the derivatives of $C_{q_3 r_1}$ and $C_{q_3 r_2}$ at $\langle z_1, y_0 \rangle$ are close to that of $C_{\hat{q}_3 \hat{r}}$. We can now find $q_1, q'_1 \in Q'_1$ such that $C_{q_3 r_1 q_1}$ and $C_{q_3 r_2 q'_1}$ have the same derivative at p_0 as $X = C_{\hat{q}_3 \hat{r} \hat{q}_1}$ does and hence also C_{q_0} (this is possible by Lemma 1.18 and continuity arguments in $\mathcal{M}$). By Lemma 1.21, the curve C_{q_0} touches $C_{q_3 r_1 q_1}$ and $C_{q_3 r_2 q'_1}$ at p_0 . It follows from (*) that s_1 belongs to both $C_{q_3 r_1 q_1}$ and $C_{q_3 r_2 q'_1}$. It is not hard to see that q_1 and q'_1 are each $\mathcal{M}$ -generic over r_1, r_2, r_3, r_4, q_0 (because each of them is inter-definable with q_3 over the r_i and q_0). For the same reason, we can find $q'_3 \in Q'_1$ such that the derivative of $C_{q'_3 r_3 q_1}$ at p_0 equals to that of C_{q_0} .

Also, by our assumption on the r_i , the curves

$$C_{q_1}^{-1} C_{r_1} C_{r_2}^{-1} C_{q'_1} \quad \text{and} \quad C_{q_1}^{-1} C_{r_3} C_{r_4}^{-1} C_{q'_1}$$

have the same derivative at p_0 and therefore

$$C_{q_1}^{-1} C_{r_1} C_{q_3}^{-1} C_{q_3} C_{r_2}^{-1} C_{q'_1} \quad \text{and} \quad C_{q_1}^{-1} C_{r_3} C_{q'_3}^{-1} C_{q'_3} C_{r_4}^{-1} C_{q'_1}$$

have the same derivative at $\langle x_0, x_0 \rangle$.

Because $C_{q_3 r_1 q_1}$, $C_{q_3 r_2 q'_1}$ and $C_{q'_3 r_3 q_1}$ all have the same derivative at p_0 it follows from the above that $C_{q'_3 r_4 q'_1}$ has the same derivative as well and therefore, by Lemma 1.21, it touches C_{q_0} at p_0 . Hence $s_1 \in C_{q'_3 r_4 q'_1}$ and so $\langle r_1, r_2, r_3, r_4 \rangle \in T$.

The proof of (2). We prove: for every generic tuple $r_1, r_2, r_3 \in Q_2^3$ there are finitely many (and at least one) r_4 such that $\langle r_1, r_2, r_3, r_4 \rangle \in T$. By the strong minimality of $\mathcal{N}$ this is a first order statement. Because $Q_2 \times Q_2 \times Q_2$ has Morley degree 1, it is sufficient to prove this for any particular $\mathcal{N}$ -generic triple in Q_2^3 . This can be done similarly to (1). Fix $\langle r_1, r_2, r_3 \rangle \in Q'_2 \times Q'_2 \times Q'_2$, $\mathcal{M}$ -generic over A , where the r_i are close to $\hat{r}$. As in the proof of (1), there is $r_4 \in Q'_2$ such that $C_{r_1 r_2}$ and $C_{r_3 r_4}$ have the same derivative at $\langle z_1, z_1 \rangle$. Moreover, each pair r_1, r_2 and r_3, r_4 is generic, so by (1)

we have $\langle r_1, r_2, r_3, r_4 \rangle \in T$. Actually, as the argument in (1) shows, to witness this we may choose $q_3 \in Q'_3$ as generic as we wish, and then find the appropriate q_1, q'_1, q'_3 .

To finish the proof of (2), we need to show that $r_4 \in \text{acl}_{\mathcal{N}}(r_1, r_2, r_3, A)$. We first prove:

Claim $r_4 \in \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 q_3 A)$.

Indeed, once we choose q_3 , the family $Y = \{C_{q_3 r_1 q} : q \in Q_1\}$ is one-dimensional, through p_0 . Notice that $s_1 \notin \text{dcl}_{\mathcal{M}}(p_0)$ for otherwise there will be infinitely many curves from $\mathcal{F}$ going through p_0 and s_1 , contradicting the normality of $\mathcal{L}$. Since q_3, r_1 were chosen generically over A there are at most finitely many curves in the family Y which go through both p_0 and s_1 . Hence, $q_1 \in \text{acl}_{\mathcal{N}}(r_1 q_3 A)$. Similarly, $q'_1 \in \text{acl}_{\mathcal{N}}(r_2 q_3 A)$, $q'_3 \in \text{acl}_{\mathcal{N}}(r_3 q_1)$ and hence q'_3 is also in $\text{acl}_{\mathcal{N}}(r_1 r_2 r_3 q_3 A)$. Finally, we may conclude in a similar fashion that $r_4 \in \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 q_3 A)$, as needed.

Assume towards a contradiction that $r_4 \notin \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 A)$. Then, by the Exchange Principle and the claim above, $q_3 \in \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 r_4 A)$ and in particular, $q_3 \in \text{acl}_{\mathcal{M}}(r_1 r_2 r_3 r_4 A)$. This contradicts our freedom, as discussed above, to choose q_3 from Q'_3 to be $\mathcal{M}$ -generic over $r_1 r_2 r_3 r_4 A$. End of (2).

This finishes the proof of Claim 2.7, and we can now return to the task of finding a group configuration in $\mathcal{N}$.

For $r \in Q'_2$ let $d(r)$ denote the derivative of C_r at $\langle z_1, z_1 \rangle$. Notice that the derivative of $C_{r_1 r_2}$ at z_1 is $d(r_1)/d(r_2)$. It follows from the properties of T described in 2.7 that, for r_1, r_2, r_3 sufficiently close to each other in Q_2 and $\mathcal{M}$ -independent, there exists r_4 such that $d(r_1)/d(r_2) = d(r_3)/d(r_4)$ and $r_4 \in \text{acl}_{\mathcal{N}}(r_1 r_2 r_3 A)$.

It is now easy to get a group configuration in $\mathcal{N}$ using the (local) group $\{d(r) : r \in Q'_2\}$. End of Lemma 2.6. $\square$

Remark The assumptions of Lemma 2.6 are not vacuous. Assume for example that at every $q \in Q$, every possible derivative is represented by two different ℓ_p through q . Given any definable curve $Y \subseteq Q$, consider the curve $X = \tau(Y) \subseteq M^2$. Because every $q \in Y$ has two different $\mathcal{L}$ -tangent curves ℓ_p and $\ell_{p'}$, the duality Theorem implies that C_q is tangent to X at both p and p' . We thus have p' special for (q, X) and $p' \in \text{acl}_{\mathcal{M}}(p, q)$.

We now return to The Special Case, with X, p_0 and q_0 as before. Namely, one of the special points for (q_0, X) is in the $\mathcal{M}$ -definable closure of p_0, q_0 (over p_1, p_2, p_3). Then, by Lemma 2.6 we can interpret a 1-dimensional group in $\mathcal{N}$. Because of strong minimality, the group can be assumed to live on $\mathcal{N}$, at least locally. Namely, we may assume now that we have a local 1-dimensional abelian group operation, denoted $+$, living on N .

We now start the process all over again and consider the two families of curves obtained by compositions $C_{q_3} C_{q_2}^{-1} C_{q_1}$ and addition (using the group structure) $C_{q_3} - C_{q_2} + C_{q_3}$ of the original C_q . We may assume that we are in the case of Lemma 2.6 for both families of curves, and then we proceed by defining tangency at $\langle z_1, z_1 \rangle$ for curves of the form $C_{r_1} C_{r_2}^{-1}$ and of the form $C_{r_1} - C_{r_2}$ (subtraction in the sense of the

group we just defined). Using the underlying real closed field structure we can now obtain a definable 1-dimensional field in $\mathcal{N}$, contradicting strong minimality. $\square$

3. THE TRICHOTOMY THEOREM

Let $\mathcal{N}$ be a structure definable in an o-minimal structure $\mathcal{M}$ (with a fixed interpretation of $\mathcal{N}$ in $\mathcal{M}$). A type p in the structure $\mathcal{N}$ is called *one dimensional* if it contains an $\mathcal{N}$ -formula whose $\mathcal{M}$ -dimension is one.

Notice that the notion of dimension depends on the particular interpretation of $\mathcal{N}$ in $\mathcal{M}$. E.g., if $\mathcal{N}$ is a dense linear ordering then it can be interpreted in $\mathcal{M}$ as a subset of M , or as a subset of M^2 , with the lexicographic ordering. In the first case, every nonalgebraic 1-type in $\mathcal{N}$ is one dimensional while in the second case, there are two dimensional types, but if one fixes two points in the same fibre and then the interval between them gives rise to one-dimensional types.

If p is a one-dimensional type in $\mathcal{N}$ then the restriction of $\text{acl}_{\mathcal{N}}(\cdot)$ to the realization of p forms a pre-geometry (Exchange holds because it is true in $\mathcal{M}$).

We will need the following from [17]:

Definition 3.1. Let $\mathcal{N}$ be a dependent theory (i.e. a theory with NIP). A definable set X is *unstable in $\mathcal{N}$* if there exists an $\mathcal{N}$ -formula $\varphi(x, y)$ defined over N and $\langle a_i, b_i \rangle_{i \in \omega}$, $a_i, b_i \in X$, such that $N \models \varphi(a_i, b_j)$ iff $i < j$. If there is no such formula φ then X is said to be *stable in $\mathcal{N}$* .

By Corollary 2.6 of [17], if X is stable set in a structure with NIP then it is stably embedded, namely every N -definable subset of X^k is definable with parameters from X .

Our main tools in the proof will be the Trichotomy Theorem for o-minimal structures [20] and the the following result from [9]:

Theorem 3.2. *Let $\mathcal{N}$ be a 1-dimensional structure definable in an o-minimal structure $\mathcal{M}$. For any unstable $X \subseteq N$ there exists $\mathcal{N}$ -definable $X_0 \subseteq N$ and equivalence relation E with finite classes such that X_0/E with all its induced $\mathcal{N}$ -structure is o-minimal. In particular X_0/E is linearly ordered.*

With this in hand we can now prove:

Theorem 3. *Assume that $\mathcal{N}$ is a definable structure in an o-minimal $\mathcal{M}$ and that p is a complete 1- $\mathcal{M}$ -dimensional $\mathcal{N}$ -type over a model $\mathcal{N}_0 \subseteq \mathcal{N}$. Then:*

- (1) p is trivial (with respect to $\text{acl}_{\mathcal{N}}$).
Or, there exists an $\mathcal{N}$ -definable equivalence relation E with finite classes such that one of the following holds:
- (2) p is linear, in which case either
 - (a) p/E is a generic type of a strongly minimal $\mathcal{N}$ -definable one-dimensional group G , and the structure which $\mathcal{N}$ induces on G is locally modular. In particular, p is strongly minimal. Or,
 - (b) for every $a \models p$ there exists an $\mathcal{N}$ -definable ordered group-interval I containing a/E . The structure which $\mathcal{N}$ induces on I is a reduct of an ordered vector space over an ordered division ring.
- (3) p is rich: For every $a \models p$ there exists an $\mathcal{N}$ -definable real closed field R containing a and the structure which $\mathcal{N}$ induces on R is o-minimal.

Proof. Because p is one-dimensional it is contained in an $\mathcal{N}$ -definable one dimensional set X .

Case 1: X is a stable set in $\mathcal{N}$.

It follows (see earlier discussion) that X is stably embedded in $\mathcal{N}$. In this case, we may replace $\mathcal{N}$ by X and assume that $\mathcal{N}$ is one-dimensional and $U(\mathcal{N}) = 1$. By Theorem 2.2, $\mathcal{N}$ is necessarily 1-based. It now follows from [10] that either p is trivial or p/E is the generic type of an $\mathcal{N}$ -definable locally-modular group G , for E an $\mathcal{N}$ -definable finite equivalence relation.

Because p is one-dimensional, we must have $\dim_{\mathcal{M}}(G) = 1$ (p is generic in G , so every definable subset of G containing p is group-generic in G , namely, finitely many group-translates of it cover G . Since $\dim p = 1$ we can find such a set which is 1-dimensional). It is left to show that G is strongly minimal. Indeed, if not then some infinite co-infinite set $Y \subseteq G$ is $\mathcal{N}$ -definable. By translating Y to $0 \in G$, we may assume that 0 lies on the boundaries of both an infinite component of Y and an infinite component of $G \setminus Y$. It is not difficult to see now that the relation $x - y \in Y$ defines an ordering on a small neighbourhood of 0 , contradicting stability.

Case 2: p is nontrivial and every $\mathcal{N}$ -definable one-dimensional set containing p is unstable.

By the non-triviality of p , we can then find an $\mathcal{N}$ -definable ternary relation $R \subseteq X^3$ (definable possibly over parameters outside of N_0), witnessing the non triviality of p , such that for every independent $a, b \models p$ there are finitely many c such that $\models R(a, b, c)$, and at least one of those c is in p . Moreover, we may assume that the projection map of R on any of its two coordinates is finite-to-one, and that the image of this projection has dimension 2. By compactness, we may shrink X further so that the projection of R onto the first two coordinates equals X^2 modulo possibly a definable set of dimension one.

Because X is unstable we may apply Theorem 3.2 and find an $\mathcal{N}$ -definable infinite set $X_0 \subseteq X$ and an $\mathcal{N}$ -definable finite equivalence relation E on X_0 such that the structure which $\mathcal{N}$ induces on X_0/E is o-minimal. In order to consider it as a structure on its own right we need stable embeddedness.

By Lemma 2.3 in [20], every closed interval in an o-minimal structure is stably embedded. However, the very same proof there gives a stronger result: If $\mathcal{M}'$ is an o-minimal structure interpretable in a geometric structure $\mathcal{N}$, with $\dim_{\mathcal{N}}(M') = 1$, then every closed interval in $\mathcal{M}'$ is stably embedded in $\mathcal{N}$.

By replacing X_0 by a smaller $\mathcal{N}$ -definable set we may assume that X_0/E is a closed interval in an $\mathcal{N}$ -definable o-minimal structure, hence stably embedded in $\mathcal{N}$. Let $\mathcal{X}_0$ denote the induced structure on X_0/E .

Our goal now is to transfer the o-minimal structure from X_0/E to elements realizing p (note that a-priori we do not know that X_0 contains p). We start with $a \models p$ and $b \in X_0$ independent from a . By our assumptions on R , there exists c such that $R(a, b, c)$ and any two of $\{a, b, c\}$ are inter-algebraic over the third one. We fix c and consider the relation $\bar{R} = R(x, y, c)$.

The projection of $\bar{R}$ on the first and second coordinate is finite-to-one and $\bar{R}$ defines a finite-to-finite correspondence between $p' = tp(a/c)$ and $tp(b/c)$. The map α which sends $\langle x, y \rangle \in \bar{R}$ to y/E is finite-to-one from $\bar{R}$ onto an infinite definable subset of $\mathcal{X}_0$. Let β be the projection map of $\bar{R}$ onto the x -coordinate and for $x \in \beta\alpha^{-1}(X_0)$, let $\gamma(x) = \min_{\mathcal{X}_0}(\alpha\beta^{-1}(x))$ (the set on the right is finite hence it has a minimum in the structure $\mathcal{X}_0$).

γ is a finite-to-one map from an infinite definable set Y containing p' onto an infinite definable set Z (which we may assume to be a closed interval) in $\mathcal{X}_0$. We thus obtain a definable bijection between $Y/\sim$, for an $\mathcal{N}$ -definable finite equivalence relation $\sim$, and an infinite closed interval in $\mathcal{X}_0$. This bijection allows us to transfer the structure which $\mathcal{N}$ induces on Z onto $Y/\sim$, hence the structure which $\mathcal{N}$ induces on $Y/\sim$ is o-minimal (and stably embedded as well), call it $\mathcal{Y}$. The type $p'/\sim$ is generic in $\mathcal{Y}$.

By the Trichotomy Theorem for o-minimal structures, we may assume (after possibly shrinking Y further) that the structure of $\mathcal{Y}$ is either trivial, an ordered reduct of a group-interval in an ordered vector-space, or an o-minimal expansion of a real closed field.

The non-triviality of p easily implies the non-triviality of p' and therefore of $\mathcal{Y}$: Fix $a_1, a_2 \models p'$ independent and d such that $\langle a_1, a_2, d \rangle \in R$, and define

$$\langle x_1, x_2, x_3 \rangle \in R' \Leftrightarrow \exists y \langle x_1, x_2, y \rangle \in R \& \langle x_3, d, y \rangle \in R.$$

The relation R' witnesses the non-triviality of p' .

If p is linear then, by definition, so is p' and therefore $\mathcal{Y}$ cannot define a real closed field (the pull-back of the family of affine lines will contradict linearity) hence $\mathcal{Y}$ is an ordered group interval. If p is rich then there is a definable almost normal family of curves such that $\langle a, a \rangle$ belongs to infinitely many of them (see the discussion following the definition of a rich type). In that case $\mathcal{Y}$ must be an o-minimal expansion of a real closed field. $\square$

We conclude by pointing out that the statement of the theorem in the unstable case cannot be strengthened so that the whole of p is contained in a single o-minimal structure: Consider N the expansion of the (unordered) group of the reals by a predicate for the unit interval. In this theory the ordering is definable on every interval of finite length. Take the type $q = \{x > r : r \in \mathbb{R}\}$. It is not contained in any definable ordering, but after fixing $b \models q$ the set $b < x < b + 1$ is contained in a definable o-minimal structure.

4. APPENDIX A: A PROOF OF V. D. DRIES' THEOREM ON LIMIT SETS

In this section we give we give a proof of Theorem 1.27. The proof is due to v.d. Dries, and relies in parts on [4], with minor changes intended to make it more self-contained.

We restate the theorem:

Theorem 4.1. *Let $\mathcal{M}$ be an o-minimal expansion of a field, $\mathcal{S}$ a definable family of definable closed subsets of M^k , parameterised by Z . Let $\tilde{\mathcal{S}} := \{\mathcal{S}(\gamma) \mid \gamma : (0, 1) \rightarrow Z, \text{ definable}\}$. Then $\tilde{\mathcal{S}}$ is $\mathcal{M}$ -definable and $\dim(\tilde{\mathcal{S}} \setminus \mathcal{S}) < \dim \mathcal{S}$.*

Because of the presence of field, we may assume that all curves in Z are given by $\gamma : (0, +\infty) \rightarrow Z$. It is not hard to check, using definable choice, that the collection $\{\mathcal{S}(\gamma) \mid \gamma : (0, +\infty) \rightarrow Z\}$ depends only on the class $\{\mathcal{S}_a : a \in X\}$ and not on the parameterisation. Hence, we may assume that every set in $\mathcal{S}$ appears exactly once in the family and in particular $\dim \mathcal{S} = \dim Z$ (otherwise re-parameterise, see §3 of [4]).

The proof goes through the theory of tame elementary pairs (see §8 of [4] for the details). Recall that for an o-minimal expansion $\mathcal{N}$ of a real closed field, a

tame extension $\mathcal{N} \not\preceq \mathcal{R}$ is one in which $\mathcal{N}$ is Dedekind complete. The theory of tame elementary pairs is the theory of those structures $(\mathcal{R}, \mathcal{N}, \text{st})$ where $\mathcal{R}$ is a tame extension of $\mathcal{N}$ and $\text{st} : R \rightarrow N$ is the standard part map. Note that st is defined on V , the convex hull of N in R . To simplify the notation, whenever $Y \subseteq R$ we will write $\text{st}(Y)$ for $\text{st}(Y \cap V)$.

The crucial property of the theory of tame pairs is:

Proposition 4.2. *Suppose that $(\mathcal{R}, \mathcal{N}, \text{st}) \models T_{\text{tame}}$ and $Y \subseteq N^k$ is definable in $(\mathcal{R}, \mathcal{N}, \text{st})$, then Y is definable in $\mathcal{N}$. Moreover, the theory T_{tame} is complete modulo $\text{Th}(\mathcal{N})$.*

From this we readily get a weaker version of the theorem, one which actually suffices for our purposes:

Lemma 4.3. *Let $\mathcal{M}$ be an o-minimal expansion of a field, $\mathcal{S}$ a $\emptyset$ -definable family of definable subsets of M^k , parameterised by Z . Let $\tilde{\mathcal{S}} := \{\mathcal{S}(\gamma) : \gamma : (0, +\infty) \rightarrow Z, \text{ definable}\}$. Then $\tilde{\mathcal{S}}$ is $\mathcal{M}$ -definable and $\dim \tilde{\mathcal{S}} = \dim \mathcal{S}$.*

Proof. By replacing $\mathcal{S}$ with the family $\{\text{cl}(A) : A \in \mathcal{S}\}$ we may assume that all the sets in $\mathcal{S}$ are closed (it is easy to check that the family of limit sets does not change). We may assume that $\mathcal{M}$ is saturated enough to assure that every M -definable set has (in M) a generic point. Let $\tau > M$ and $N = M\langle\tau\rangle$, the prime model over $M \cup \{\tau\}$. Then M is Dedekind complete in N and $(\mathcal{N}, \mathcal{M}, \text{st}) \models T_{\text{tame}}$. Let $\gamma : (0, +\infty) \rightarrow Z$ be any curve definable over M . It is not hard to verify that $\mathcal{S}(\gamma) = \text{st} \mathcal{S}_{\gamma(\tau)}$. By Proposition 4.2, and by compactness, the definable family $\{\text{st} \mathcal{S}_a : a \in Z(N)\}$ is $\mathcal{M}$ -definable. Since every $a \in Z(N)$ is of the form $\gamma(\tau)$ for some $\mathcal{M}$ -definable curve γ , the definability of $\tilde{\mathcal{S}}$ in $\mathcal{M}$ follows. Because every automorphism of $\mathcal{M}$ leaves $\mathcal{S}$ invariant and therefore also $\tilde{\mathcal{S}}$, it follows that $\tilde{\mathcal{S}}$ is actually $\emptyset$ -definable.

Now let Z_1 be a parameter set for $\tilde{\mathcal{S}}$ (so Z_1 is $\emptyset$ -definable in $\mathcal{M}$). Again, we assume that every set in $\tilde{\mathcal{S}}$ is represented exactly once in the family. Let $b \in Z_1(M)$ be generic over $\emptyset$. There is some $\gamma : (0, +\infty) \rightarrow Z$ such that $\tilde{\mathcal{S}}_b = \mathcal{S}(\gamma) = \text{st} \mathcal{S}_{\gamma(\tau)}$. If $\gamma(\tau) \in M^k$ then γ is eventually constant and hence $\mathcal{S}_\gamma \in \mathcal{S}$. Assume then that $\gamma(\tau) \notin M$ and let $\mathcal{P} = \langle\gamma(\tau)\rangle$, the model generated by $\gamma(\tau)$ over $\emptyset$. Consider the tame pair $(\mathcal{P}, \text{st} \mathcal{P}, \text{st})$. It follows from the completeness part of Proposition 4.2 that $\text{st} \mathcal{S}_{\gamma(\tau)} = \tilde{\mathcal{S}}_{b'}$ for some $b' \in Z_1(\text{st} \mathcal{P})$. By our assumption, we must have $b' = b$.

Since $\gamma(\tau) \in Z$, we have $\dim(\gamma(\tau)/\emptyset) \leq \dim(\mathcal{S})$, and therefore

$$\dim(\mathcal{S}) \geq \dim(\mathcal{P}/\emptyset) \geq \dim(\text{st} \mathcal{P}/\emptyset).$$

We conclude that $\dim(b/\emptyset) \leq \dim(\mathcal{S})$ therefore $\dim(\tilde{\mathcal{S}}) = \dim(Z_1) \leq \dim(\mathcal{S})$. $\square$

As mentioned above, the previous lemma suffices for our purposes. For the sake of completeness, however, we give the proof of v.d. Dries' theorem in its full strength.

The main part of the proof is the following:

Lemma 4.4. *For $\mathcal{S}$ as above, assume that all sets in $\mathcal{S}$ are closed and $\emptyset \notin \mathcal{S}$. Let $M \prec N \prec R$, be such that M, N are Dedekind complete in R , and let $\text{st} : R \rightarrow N$ be the associated standard part map, V the convex hull of N in R . Let $a \in Z(R) \cap V^n$ be such that $M\langle a \rangle \subseteq V$. Then $\text{st} a \in Z(N)$ and $\text{st}(\mathcal{S}_a(R)) = \mathcal{S}_{\text{st} a}(N)$.*

Proof of the lemma. Since $M\langle a \rangle \subseteq V$ the map $\text{st}|_{M\langle a \rangle} : M\langle a \rangle \rightarrow N$ is elementary. So $\text{st} a \in Z(N)$.

Consider the function $f : Z \times M^n \rightarrow M$ given by $f(z, y) = d(\mathcal{S}_z, y)$ (with $d(z, y) = \sqrt{z^2 + y^2}$ and $d(S, y) = \inf\{d(z, y) : z \in S\}$). So for each $z \in Z$ the function $f_z(y) : f(z, y)$ is continuous. Hence by §6 (Section 2) of [3] there is a partition of Z into definable subsets $Z_1, \dots, Z_k$ such that each restriction $f_i|_{X_i \times M^n}$ is continuous. Restricting Z , this allows us to assume that f is continuous on Z . Let $b \in \mathcal{S}_a(R) \cap V^n$; so $f(a, b) = 0$. Since f is continuous (and $\text{st } a \in Z$), we get that $f(\text{st } a, \text{st } b) = \text{st } f(a, b) = 0$. i.e. $\text{st } b \in \text{cl } \mathcal{S}_{\text{st } a}$. Since we assume all the $\mathcal{S}_z$ to be closed, this is what we wanted.

Conversely, if $c \in \mathcal{S}_{\text{st } a}$ we have that $\text{st } c = c$ and therefore $f(\text{st } a, \text{st } c) = 0$. Reversing the above argument, we get that $f(a, b)$ is infinitesimal. By the definition of f this means that $d(b, c)$ (computed in R) is infinitesimal for some $b \in \mathcal{S}_a(R)$. Thus, $c = \text{st } b \in \mathcal{S}_a(R)$, as desired. $\square$

The rest of the proof is a rehash to what we did in Lemma 4.3: By Proposition 4.2 the set $\tilde{\mathcal{S}} \setminus \mathcal{S}$ is $\mathcal{M}$ -definable, and denote the corresponding parameter set Z' . It's enough to show that $\dim Z' < \dim Z$. We take $\mathcal{M} \prec \mathcal{N} \prec \mathcal{R}$ as in the previous lemma, and denote $\text{st}_M : R \rightarrow M$, $\text{st} : R \rightarrow N$ the corresponding standard part maps. As in the proof of Lemma 4.3 we obtain:

$$\{\tilde{\mathcal{S}}_b : b \in Z'(M)\} = \text{st}_M \mathcal{S}_a \setminus \{\mathcal{S}_a : a \in Z(M)\}$$

Since $(\mathcal{R}, \mathcal{M}, \text{st}_M) \equiv (\mathcal{R}, \mathcal{N}, \text{st})$ we also get

$$\{\tilde{\mathcal{S}}_b : b \in Z'(N)\} = \text{st } \mathcal{S}_a \setminus \{\mathcal{S}_a : a \in Z(N)\}$$

By the previous lemma, if $b \in Z'(N)$ is generic and $a \in Z(R)$ is such that $\text{st } \mathcal{S}_a(R) = \tilde{\mathcal{S}}_b(N)$, it cannot be that $M\langle a \rangle \subseteq V$. Thus, $\dim(\text{st } M\langle a \rangle / M) < \dim(a/M)$, and since b was arbitrary, the desired result follows exactly as in the proof of Lemma 4.3.

5. APPENDIX B: THE GENERAL O-MINIMAL CASE

In this appendix we adapt the main results of this paper to the general o-minimal context (i.e. removing the assumption that $\mathcal{M}$ expands a real closed field). From the proof of the results in Section 3 it is clear that we only need concern ourselves with the proof of Theorem 2.2.

In the first part of this appendix we will prove two important technical facts for $\mathcal{N}$ strongly minimal, 1-dimensional and definable in a sufficiently saturated o-minimal structure $\mathcal{M}$:

- $\mathcal{N}$ has finitely many non-orthogonality classes (in the o-minimal sense).
- $\mathcal{M}$ has definable choice for $\mathcal{M}$ -definable subsets of N^k .

The second part of the appendix uses these two facts to overcome the main obstacle in the generalisation of 2.2: Lemma 1.31. This lemma relies heavily on v.d. Dries' Theorem on limit sets (1.27), which is only known to be true in expansions of real closed fields.

Finally, we will go systematically through all places where the existence of a field was used, and explain briefly how to avoid it.

Throughout this section $\mathcal{N}$ will be a 1-dimensional structure *definable* in a densely ordered o-minimal $\mathcal{M}$. It seems plausible that a further generalisation of the theorem

to the case where $\mathcal{N}$ is only interpretable in $\mathcal{M}$ can be achieved, but this will definitely require more work.

Claim 5.1. *Without loss of generality, N is a dense subset of M .*

Proof. Since $\dim_{\mathcal{M}} N = 1$ it is a finite union of 1-cells and 0-cells (in M^k for some k). Note that removing finitely many points cannot alter the truth of Theorem 2.2 and therefore neither of Theorem 2 and Theorem 1. Hence we may assume that N is given by a union of 1-cells only. Each 1-cell is definably homeomorphic to an interval in M and therefore inherits an ordering. Thus each 1-cell, with its induced structure from $\mathcal{M}$ can be viewed as an o-minimal structure on its own. We now take the disjoint union of the 1-cells of N , ordering them together arbitrarily, so that each cell is an open interval in this ordering. We add endpoints between these intervals and obtain an o-minimal structure $\mathcal{M}'$, such that N , without its 0-cells, is a definable dense subset of $\mathcal{M}'$. We can now replace $\mathcal{M}$ by $\mathcal{M}'$. $\square$

5.1. Non-orthogonality and definable choice. Since we can no longer assume that $\mathcal{M}$ has elimination of imaginaries, and as the work throughout the paper was carried out in $\mathcal{M}$ (and not in $\mathcal{M}^{eq}$) we should be careful when using $\mathcal{N}$ -interpretable subsets - as these will not, a priori, be definable in $\mathcal{M}$. Recall:

Definition 5.2. A theory T has *weak elimination of imaginaries* if for every $M \models T$ and $e \in M^{eq}$ there are $a_1, \dots, a_n \in M$ such that $e \in \text{dcl}(a_1, \dots, a_n)$ and $a_i \in \text{acl}(e)$ for all $1 \leq i \leq n$.

It is fairly easy to check (See, e.g., [12]) that:

Lemma 5.3. *Assume T is strongly minimal and $\text{acl}(\emptyset)$ is infinite then T has weak elimination of imaginaries.*

Hence, by adding constants to $\mathcal{N}$ we may assume that it has weak EI.

Definition 5.4. Let $\mathcal{M}$ be an o-minimal structure. $a_1, a_2 \in M$ are *non-orthogonal* if there exists a continuous monotone definable bijection $f : I_{a_1} \rightarrow I_{a_2}$ for some open intervals $I_{a_i} \ni a_i$ sending a_1 to a_2 (see more in [20]).

If $S \subseteq M$ is a definable set and $a \in M$ any element, we say that $b \not\perp a$ for every $b \in S$, *uniformly in a* , if there exists a definable family of definable functions $F : M^2 \rightarrow M$ such that for all $x \in S$ the function $f_x(y) := F(x, y)$ witnesses that $x \not\perp a$.

The notion of (non) orthogonality plays an important role in following arguments. The key observation, which makes everything else work is:

Lemma 5.5. *If $\mathcal{N}$ is non-locally modular then there exists $a \in \mathcal{N}$ such that $a \not\perp b$ uniformly in b (in the o-minimal sense, obviously) for all $b \in M$ outside a finite set.*

Proof. Choose $a \in \mathcal{N}$ generic (in the o-minimal sense). By weak EI we can find $\mathcal{F}$ an almost normal 2-dimensional family of $\mathcal{N}$ -definable plane curves, as provided by non local modularity (see Fact 2.4). Hence for $b \in \mathcal{N}$, o-minimally generic over a , the point $\langle a, b \rangle$ is generic on a generic $\mathcal{F}$ -curve through $\langle a, b \rangle$, implying that such a curve is, locally near $\langle a, b \rangle$, the graph of a continuous monotone function with $f(a) = b$, so $a \not\perp b$.

Consider now the set S of all $b \in \mathcal{N}$ such that there is no $\mathcal{F}$ -line through $\langle a, b \rangle$ which is locally near $\langle a, b \rangle$ the graph of a continuous monotone function. This is an o-minimally definable set, and by what we have just shown, it must be finite. $\square$

Remark 5.6. The above lemma seems to remain true under the weaker assumption that $\mathcal{N}$ is non-trivial. In that case, if $\mathcal{N}$ is locally modular, use the fact that there exists an $\mathcal{N}$ -definable equivalence relation E with finite classes such that N/E is a co-finite set in a strongly minimal abelian group.

The following is easy:

Lemma 5.7. *If $\mathcal{M}$ has finitely many non-orthogonality classes then $\mathcal{M}$ does not have a generic trivial type.*

Proof. Assume that $e \in M$ is a generic trivial point. Let $I \ni e$ be a closed interval such that every point in I is trivial (see, e.g., the introduction to [20]). We claim that every $a, b \in I$ that are independent must be orthogonal to each other. Indeed, if not then there is a definable, continuous, strictly monotone function f sending a to b . It follows that for every b' near b there is, uniformly in b' , a definable $f_{ab'}$ witnessing the non-orthogonality of a and b' . It follows that b is nonorthogonal, uniformly, to every element in some neighbourhood of it. It is not difficult to see that this contradicts the triviality of b . $\square$

We can now prove that $\mathcal{M}$ has definable choice:

Lemma 5.8. *If $\mathcal{N}$ is strongly minimal and non-trivial (and we still assume that N is a dense subset of M) then $\mathcal{M}$ has definable Skolem functions. I.e. for every $\mathcal{M}$ -definable formula $\varphi(\bar{x}, \bar{y})$ there is an $\mathcal{M}$ -definable function f_φ such that $\exists \bar{y} \varphi(\bar{x}, \bar{y}) \rightarrow \varphi(\bar{x}, f_\varphi(\bar{x}))$.*

Proof. Since $M \setminus N$ is finite, it is enough to consider definable subsets of N^k . By cell decomposition, we only need to consider cells, and therefore, it suffices to prove that given a z -definable interval I_z we can find a z -definable point in I_z .

If $\mathcal{N}$ is locally modular, then since it is non-trivial there are an $\mathcal{N}$ -definable vector space V , a finite equivalence relation E on N and a definable injection $f : N/E \rightarrow V$ (see, [25]) whose image is co-finite in V . By [6], $\mathcal{M}$ has definable choice for definable subsets of V , this gives an element $z_0 \in f(I)$ and we take $z = \min\{f^{-1}(z_0)\} \in I$.

So we may assume that $\mathcal{N}$ is not locally modular. Let $(c, d) \subseteq N$ be any interval. We show that we can choose a point $e \in (c, d)$ uniformly in c, d . Combining the previous lemma with Lemma 5.7 for all but finitely $x \in N$ there exists, uniformly in x , an interval I_x on which a group-interval (not a group!) is defined. If c, d are not both of the finitely many exceptional points, there is (without loss) an interval $I_c \ni c$ as above. If $d \notin cl(I_c)$ then setting $e := \sup I_c \in (c, d)$ we are done. Otherwise $(c, d) \subseteq I_c$ and we can set $e = (c +_c d)/_c 2_c$ (in the sense of the local group defined on I_c) if $d < \sup I_c$ or $c +_c 1_c$ otherwise. Since the number of exceptional points, and hence the number of intervals with exceptional end-points, is finite, this gives us definable choice for sub intervals in N . $\square$

5.2. Limit sets in structures without fields. Our next task is to slightly refine the notion of non-orthogonality in o-minimal structures.

Definition 5.9. Let $\mathcal{M}$ be an o-minimal (saturated enough) structure, $a, b \in M \cup \{\pm\infty\}$. We say that $a^+ \not\perp b^+$ if there exist intervals $I_a^+ := (a, a_\epsilon)$, $I_b^+ := (b, b_\epsilon)$

and a definable (with parameters) continuous monotone bijection $f : I_a^+ \rightarrow I_b^+$ (so $\lim_{t \rightarrow a} f(t) = b$). For $\varepsilon_i \in \{+, -\}$ we define analogously, $a^{\varepsilon_1} \not\perp b^{\varepsilon_2}$ and $a^\varepsilon \not\perp \pm\infty$.

Consider the structure $\mathcal{M}^\pm$ consisting of $\mathcal{M}$ with all its structure, as well as a new sort $M^\pm = (M \cup \{\pm\infty\}) \times \{+, -\}$ and the projection $\pi : M^\pm \rightarrow M \cup \{\pm\infty\}$. The relation $a^{\varepsilon_1} \not\perp b^{\varepsilon_2}$ is an equivalence relation on the elements of $M \times \{+, -\}$, so we can define:

Definition 5.10. A complete set of representatives of the non-orthogonality classes of $\mathcal{M}^\pm$ is a subset of $\mathcal{M}^\pm$ in which every nonorthogonality class is represented.

Definition 5.11. Assume that $\mathcal{M}$ is o-minimal and every a, b outside a finite set F are non-orthogonal to each other, uniformly in a, b . For the purposes of this note, we call such $\mathcal{M}$ *uniformly unidimensional*.

Lemma 5.12. *Let $\mathcal{M}$ be uniformly unidimensional, then $\mathcal{M}$ has at most finitely many trivial points.*

Proof. This is immediate from 5.7. □

Note that by the previous lemma and (the proof of) Lemma 5.8 a uniformly unidimensional structure $\mathcal{M}$ has definable choice. Moreover, uniformly unidimensional structures have curve selection for non-degenerate points. More precisely, if $S \subseteq M^r$ is a definable set and $q := (q_1, \dots, q_r) \in \partial S$ is such that $q_i \notin F$, then there are $a < b \in M$ and a definable curve $\gamma : (a, b) \rightarrow S$ such that $\lim_{t \rightarrow a} \gamma(t) = q$.

Definition 5.13. Let $\mathcal{M}$ be uniformly unidimensional. $a \in M$ is *degenerate* if one of a^+, a^- is not in the generic non-orthogonality class of $\mathcal{M}$.

A few clarifications may be in place:

Remark

(1) If $\mathcal{M}$ has finitely many non-orthogonality classes so does $\mathcal{M}^\pm$. The converse is true as well: Let $\{a_1^{\varepsilon_1}, \dots, a_k^{\varepsilon_k}\}$ be a set of representatives of the non-orthogonality classes of $\mathcal{M}^\pm$. For $b \in \mathcal{M}$ define the non-orthogonality type of b to be the pair (i, k) ($1 \leq i, j \leq k$) such that $b^+ \not\perp a_i^{\varepsilon_i}$ and $b^- \not\perp a_j^{\varepsilon_j}$. It will be enough to show that if b, d have the same non-orthogonality type, they are non-orthogonal in the usual sense, which is a simple exercise in concatenation and composition of monotone functions.

(2) The number of non-orthogonality classes, when finite, depends only on $Th(\mathcal{M})$. Indeed, assume that $\{a_1^{\varepsilon_1}, \dots, a_k^{\varepsilon_k}\}$ is a complete set of representatives of the non-orthogonality classes of $\mathcal{M}$. Because of the saturation of $\mathcal{M}$ the type stating that $x^+ \in M$ is orthogonal to each of the $a_i^{\varepsilon_i}$ is inconsistent. It follows that, uniformly in a , every a^+ is nonorthogonal to one of the $a_i^{\varepsilon_i}$, and similarly for a^- . By quantifying over the a_i we get in every model $\mathcal{M}_0$ of $Th(\mathcal{M})$ k -many non-orthogonality classes in $\mathcal{M}_0^\pm$.

(3) It is possible that a^{ε_1} and b^{ε_2} are non-orthogonal to each other, while a is a degenerate point and b is a non-degenerate point. For example, consider $(\mathbb{R}; <, + | (\mathbb{R}_{>0})^2)$. We have $0^+ \not\perp r$ for every positive r but 0 is a degenerate point.

Fix $\mathcal{M}$, a uniformly unidimensional structure.

We now start our treatment of limit sets in $\mathcal{M}$. Let $F^\varepsilon = \{a_1^{\varepsilon_1}, \dots, a_k^{\varepsilon_k}\}$ be the set of representatives of non-orthogonality classes in $\mathcal{M}^\pm$ which are orthogonal to c^+ for

a fixed generic c . In particular, $F^\epsilon \cup \{c^+\}$ is a complete set of such representatives. We let $F = \{a_1, \dots, a_k\}$. Clearly, $F \subseteq \text{dcl}_M(\emptyset)$.

Recall that in the case of o-minimal expansions of fields, given a definable family of definable sets $\mathcal{S}$, with parameter set Z_1 , the limit family $\mathcal{S}$ was defined as $\{\mathcal{S}(\gamma) : \gamma : (0, 1) \rightarrow Z_1 \text{ definable}\}$. In the presence of more than one non-orthogonality class not all definable curves γ with image in Z_1 can have the same domain. In uniformly unidimensional structures, we handle separately limits along curves $\gamma : (a, b) \rightarrow Z_1$ with a^+ degenerate and those curves for which a^+ is non-degenerate. Clearly, we can handle similarly curves $\gamma : (b, a) \rightarrow Z_1$ and a^- .

Finally, it may happen that for some curve γ the limit set $\mathcal{S}(\gamma)$ is finite. To avoid unpleasant technicalities, and since in the application we will always be interested solely in limit sets that are infinite, we will restrict ourselves to those. We start with the former case:

Lemma 5.14. *Let $\mathcal{M}$ be as above, $\mathcal{S}$ an almost normal family of definable curves with parameter set Z_1 , and $\gamma : (a, b) \rightarrow Z_1$ a definable curve, $a^+ \in F^\epsilon$. Assume that the limit set $\mathcal{S}(\gamma)$ is infinite and let $x = \langle x_1, \dots, x_n \rangle$ be a generic element in $\mathcal{S}(\gamma)$. Then there exists some i such that $x_i \in \text{dcl}(\emptyset)$. In particular, by genericity $x'_i = x_i$ for all $x' \in \mathcal{S}(\gamma) \cap U$ for some open $U \ni x$.*

Proof. As before we may assume that all the curves in $\mathcal{S}$ are closed (after moving to the family of closures we still have an almost normal family). Assume that the lemma fails. Then for some generic $x \in \mathcal{S}(\gamma)$ all coordinates are non-orthogonal to some generic $c \in M$. Since c is not trivial, there is, by the Trichotomy Theorem, a group interval defined around c . Pulling back this group structure to x (on each coordinate separately) we may assume that x lives in an expansion of a group interval. In particular, there is a definable family of open sets $U_r \ni x$, with $r \in M$ such that $\lim_{r \rightarrow c} U_r = \{x\}$ (the existence of this family is our only use of the group structure near x). Since $x \in \mathcal{S}(\gamma)$, for every r there exists $t(r) \in (a, b)$ such that $S_{\gamma(t(r))} \cap U_r \neq \emptyset$. Since $a^+ \in F$, it cannot be that $t(r) \rightarrow a$ as $r \rightarrow c$. Therefore, $\lim_{r \rightarrow c} t(r) = a_0 > a$. It follows that $x \in S_{\gamma(a_0)}$. But this is impossible: we can repeat the process, finding $a < \dots < a_n < \dots < a_1 < a_0$ with $x \in S_{\gamma(t(a_i))}$ for all i , i.e. x is in infinitely many of the $S_{\gamma(t)}$, so without loss, it appears in all of them (because this is a 1-dimensional family). Since we can repeat this for infinitely many x 's in $\mathcal{S}_\gamma$ - this would contradict the normality of $\mathcal{S}$. $\square$

For $\gamma : (a, b) \rightarrow M^2$ a definable curve, we say that γ is of rank k over A if the image of the interval (a, b) under γ is a curve of rank k over A .

From now on we also assume: **Every generic type in $\mathcal{M}$ is rich, namely it is contained in an $\mathcal{M}$ -definable real closed field.**

Lemma 5.15. *Let $\mathcal{M}$ be as above. Let $\gamma = \langle \gamma_1, \gamma_2 \rangle : (a, b) \rightarrow M^2$ be a definable curve, with $a^+ \notin F^\epsilon$. If γ is of rank $k \geq 3$ over A then there are open intervals $I_1, I_2 \subseteq M$ such that:*

- (i) *For some $a < b_1 < b$, the curve $\gamma(a, b_1)$ is contained in $I_1 \times I_2$;*
- (ii) *I_1, I_2 are definable over $B \supseteq A$ and the rank of $\gamma|(a, b_1)$ over B is still k .*
- (iii) *I_1, I_2 are definably isomorphic to some open intervals in a definable real closed field R .*

Proof. Since γ has rank k over A , there exists an A -definable almost normal family of curves $\mathcal{X} = \{X_d : d \in D\}$, such that $\gamma(a, b) = X_d$ for some generic $d \in D$, and

$\dim(D) = k$. We first note that for every generic $x = \langle x_1, x_2 \rangle \in X_d$, we have $x_1, x_2 \notin \text{dcl}(\emptyset)$. Indeed, otherwise X_d would be locally the graph of a constant function and hence, in some M^2 -neighbourhood of x , the family $\mathcal{X}$ will be 1-dimensional, contradicting the fact that $\mathcal{X}$ is almost normal and of rank at least 3.

Let $\langle p_1, p_2 \rangle \in (M \cup \{\pm\infty\})^2$ be the limit of $\gamma(t)$ as t tends to a . Because of our last remark, the functions γ_1, γ_2 are nowhere constant hence for some $\epsilon_1, \epsilon_2 \in \{\pm\}$ we have $p_1^{\epsilon_1} = \lim_{t \rightarrow a} \gamma_1(t)$ and $p_2^{\epsilon_2} = \lim_{t \rightarrow a} \gamma_2(t)$ (by $p^+(p^-) = \lim_{t \rightarrow a} \gamma_1(t)$ we mean that $\gamma(t)$ tends to p from above (below)). To simplify notation, let us assume that $\epsilon_1 = \epsilon_2 = +$. Since $a^+ \notin F^c$, it is nonorthogonal to c^+ , for our fixed generic c . Let R denote a fixed definable real closed field such that $c \in R$.

There are now several cases to consider:

Case 1 $p_1, p_2 \in \text{dcl}(\emptyset)$.

We consider intervals $I_1 = (p_1, p'_1)$, $I_2 = (p_2, p'_2)$, for p'_1, p'_2 generic and independent from d over A . By choosing b_1 sufficiently small, we have $\gamma(a, b_1) \subseteq I_1 \times I_2$. Because γ_1, γ_2 witness the fact that $p_1^+, p_2^+ \not\perp c^+$, I_1, I_2 are definably isomorphic to open intervals in the field R .

Case 2 $p_1, p_2 \notin \text{dcl}(\emptyset)$.

In this case, p_1 and p_2 are nonorthogonal to c and therefore each is contained in a definable real closed field. We can then find open intervals $I_1 \ni p_1, I_2 \ni p_2$, definable over generic parameters which are independent from d over A , such that each I_i is contained in a definable real closed field. Because all generic points are uniformly non-orthogonal to each other, the two intervals are definably isomorphic to intervals in R .

Case 3 $p_1 \in \text{dcl}(\emptyset)$ while $p_2 \notin \text{dcl}(\emptyset)$.

In this case, we take $I_1 = (p_1, p'_1)$ and I_2 an interval containing p_2 and proceed as before. We handle similarly the symmetric fourth case. $\square$

Lemma 5.16. *Let $\mathcal{M}$ be as above. Let $\mathcal{S}$ be a definable two-dimensional almost normal family of curves, all contained in a two dimensional set Q , and having a parameter set $P \subseteq M^2$. Let $\gamma : (a, b) \rightarrow P$ be a definable curve of rank $k \geq 3$.*

Assume that S_γ is infinite. If q is a generic element in $\mathcal{S}(\gamma)$, with $\dim(q) = 2$ then there exists an open neighbourhood U of q such that the rank of $\mathcal{S}(\gamma) \cap U$ is at most 2.

Proof. By Lemma 5.14, $a^+ \notin F^c$, for otherwise we will have $\dim(q) \leq 1$.

If $p = \lim_{t \rightarrow a} \gamma(t)$ then, by Lemma 5.15, there exists $b_1 \in (a, b)$ and an open rectangular box $V = I_1 \times I_2$ such that $\gamma(a, b_1) \subseteq V$, and V is definably isomorphic to an open rectangular box in a definable real closed field R . Moreover, the rank of $\gamma|(a, b_1)$ over all parameters is still $k \geq 3$. Assume that $\gamma(a, b_1) = X_d$ for X in a definable family of curves $\mathcal{X} = \{X_{d'} : d' \in D\}$. Then, after possibly shrinking D (but still with $\dim(D) = k$, we may assume that for every $d' \in D$ the curve $X_{d'}$ is contained in V .

Because $\dim(q) = 2$, we may assume that $q = \langle q_1, q_2 \rangle \in M^2$, where q_1, q_2 are nonorthogonal to c . Hence, there exists an open rectangular box $U = J_1 \times J_2 \ni q$ such that J_1, J_2 are definably isomorphic to open intervals in the field R .

We can now restrict ourselves to $\mathcal{S}(U, V) = \{\mathcal{S}_{p'} \cap U : p' \in V\}$. Because U, V are definably isomorphic to open rectangular boxes in the field R , we may assume that $\mathcal{S}(U, V)$ and every limit set of this family are definable in R . We can now apply Theorem 1.27 and conclude that the family of all possible limit sets for the family

$\mathcal{S}(U, V)$ is definable of dimension at most 2. The limit curve $\mathcal{S}(\gamma) \cap U$ belongs to this family and therefore its rank is at most 2. $\square$

Finally, we can prove Lemma 1.31 without the field assumption. We formulate it here again:

Lemma 5.17. *For $\mathcal{M}$ uniformly unidimensional, let $\mathcal{F}$ be an almost normal family of plane curves parameterised by Q , $X \subseteq P$ an A -definable curve of rank $k > 2$, $p_0 \in X$ generic and assume that for $q_0 \in Q$, the curve C_{q_0} touches X at p_0 (here we identify P locally with an open set in M^2), and in addition, $\dim(p_0, q_0/\emptyset) = 3$. If p is a special point for (q_0, X) then $\dim(p/A) = 1$. In particular, p is a generic point of X .*

Proof. As was pointed out earlier on, $p \in \text{cl}(X)$ and hence $\dim(p/A) \leq 1$. By 1.6, we may assume that in a neighbourhood U of p_0 and V of q_0 , the family $\mathcal{F}$ is nice. Consider $\tau(X \cap U)$ and recall that q_0 is generic in $\tau(X \cap U)$ over A (by the Duality Theorem).

Assume, towards a contradiction, that $\dim(p/A) = 0$. Hence, q_0 is still generic in $\tau(X \cap U)$ over Ap . As was pointed out in clause (3) of the discussion following Definition 1.28, it follows that q_0 is in the Ap -definable limit set $\mathcal{L}(X, p)$. But then, by the genericity of q_0 over pA , there exists a neighbourhood $V_1 \subseteq V$ of q_0 , such that $\tau(X \cap U) \cap V_1 = \mathcal{L}(X, p) \cap V_1$. In particular, $\mathcal{L}(X, p)$ is infinite.

The limit set $\mathcal{L}(X, p)$ can also be written as $\mathcal{L}(\gamma)$ with $\gamma : (a, b) \rightarrow P$ a curve of rank $k > 2$ over A (γ determines a branch of X which contains p in its frontier). By Lemma 5.14, there exists an open neighbourhood V' of q_0 , such that $\mathcal{L}(\gamma) \cap V'$ has rank at most 2 over A . Without loss of generality, $V' = V_1$ and therefore the rank of $\tau(X \cap U) \cap V_1$ over A is at most 2. However, since the rank of X over A is $k > 2$, then the rank of $X \cap U$ is k as well and, by Lemma 1.25, the rank of $\tau(X \cap U) \cap V_1$ is k as well (see 1.20), contradiction. $\square$

5.3. Some extra details. In this final subsection, we explain - claim by claim - how to avoid any use we made of the ambient field structure on $\mathcal{M}$. As already noted, the proof of Theorem 3 relies on results of the first sections of this paper only in the treatment of the stable case, and does not otherwise rely on $\mathcal{M}$ expanding a field. It will suffice therefore to show that Theorem 2.2 as well is true independently of the field assumption. Therefore, throughout this section we will assume that $\mathcal{N}$ is a strongly minimal non-locally modular structure *definable* in an arbitrary o-minimal structure $\mathcal{M}$.

The following lemma shows that under these (a posteriori inconsistent) assumptions fields are (locally) definable in $\mathcal{N}$:

Lemma 5.18. *Let $\mathcal{M}$ be an o-minimal structure, $\mathcal{N}$ a 1-dimensional non-locally modular strongly minimal set definable in $\mathcal{M}$ such that N is dense in M . Then $\mathcal{M}$ is uniformly unidimensional and every generic type in $\mathcal{M}$ is rich.*

Proof. We already established that $\mathcal{M}$ is uniformly unidimensional (Lemma 5.5 and Lemma 5.8).

Adding constants, we may assume that $\mathcal{N}$ has weak EI. The non-local modularity of $\mathcal{N}$ gives rise to a 2-dimensional family of plane curves in $\mathcal{N}$ (and hence in $\mathcal{M}$). Such a family cannot be definable in trivial structures or in ordered vector spaces, hence by the Trichotomy Theorem for o-minimal structures a real closed field R is definable

in $\mathcal{M}$. By the unidimensionality of $\mathcal{M}$ every generic type in $\mathcal{M}$ is nonorthogonal to a generic type in R . $\square$

From the last lemma it follows that all the results of Sections 1 and 2 which uses the field structure only in a local way transfer to the present context. Here are some examples:

- (1) In Lemma 1.17: The fact that X is a graph of a C^1 -function is meaningless without the existence of an ambient field. But since we are interested in the result only locally near a generic point $p_0 = \langle x_0, y_0 \rangle \in X$ this can be solved as follows. Since $x_0, y_0 \in N$ are generic in $\mathcal{M}$ they are non-orthogonal to each other and each has an $\mathcal{M}$ -definable real closed field containing it. By non-orthogonality there exist closed (and non-trivial) $\mathcal{M}$ -intervals $I_1 \ni x_0, I_2 \ni y_0$, which are definably homeomorphic to each other. We may assume that I_1, I_2 are definably homeomorphic to open intervals in the same field R . Hence, differentiation can be take with respect to that field structure. Note, also, that by stable embeddedness, $X \cap I_1 \times I_2$ is definable, locally near p_0 using only parameters from $I_1 \cup I_2$. Thus, the lemma clearly remains true in this new formulation.
- (2) Lemma 1.18: Again, we use the fact that x_0, y_0 are nonorthogonal to generic elements in the same real closed field.
- (3) The two lemmas discussed above are crucial to prove Theorem 1.21, where the assumptions are only meaningful in the presence of an ambient field, but whose conclusion is independent of that field. Thus, we can reformulate the theorem as follows: Let $\mathcal{F}$ be a $\emptyset$ -definable (almost) normal family of plane curves of dimension greater than 1. Let X be an A -definable plane curve of rank greater than 1, $p_0 = \langle x_0, y_0 \rangle \in X$ generic in over A . Endow $\emptyset$ -definable neighbourhoods of x_0 and y_0 with a similar field structure, as above, and let d be the derivative - with respect to that field structure - of the function associated to X at x_0 . If $p_0 \in C_{q_0}$ for $q_0 \in Q$ and if $f'_{q_0}(x_0) = d$, then C_{q_0} and X touch each other at p_0 and $\dim_{\mathcal{M}}(p_0, q_0/\emptyset) = 3$.

This covers, all occurrences of the field structure in Section 1. We now turn to Section 2.

- (1) The construction of the curve X , depends only on the $\mathcal{N}$ -structure and so causes no problem. We have to make sure, however, that there exists a curve from the family $\mathcal{F}$ which touches X at a generic point. This follows, essentially, from the version of Theorem 1.21 we just formulated above. However, an alternative approach is also possible. Note that in the construction, we only need the points z_1, z_2, x_0, y_0 to be independent generics. By choosing them all very close to each other, and restricting our attention to a small interval containing them all, we may in fact assume (using stable embeddedness, as usual) that in that part of the proof we are working in an expansion of a field.
- (2) Since we checked that the results of Section 1 go through, the proof of Theorem 2.2 in the first case, (the “general case”) needs no alteration. As for the second part (the “special case”), we note that the argument is only concerned with what is going on near (in the o-minimal sense) the point p_0 . Thus choosing z_1, z_2, x_0, y_0 as suggested above, will assure that the argument will need no alteration.

REFERENCES

- [1] Elisabeth Bouscaren. The group configuration—after E. Hrushovski. In *The model theory of groups (Notre Dame, IN, 1985–1987)*, volume 11 of *Notre Dame Math. Lectures*, pages 199–209. Univ. Notre Dame Press, Notre Dame, IN, 1989.
- [2] Egbert Brieskorn and Horst Knörrer. *Plane algebraic curves*. Birkhäuser Verlag, Basel, 1986. Translated from the German by John Stillwell.
- [3] Lou Van Den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.
- [4] Lou Van Den Dries. Limit sets in o-minimal structures. In *Proceedings of RAAG summer school, Lisbon 2003*, pages 159–215, 2005.
- [5] Lou Van Den Dries. A remark on limit sets. 2006.
- [6] Mário J. Edmundo. Solvable groups definable in o-minimal structures. *J. Pure Appl. Algebra*, 185(1-3):103–145, 2003.
- [7] Jerry Gagelman. Stability in geometric theories. *Ann. Pure Appl. Logic*, 132(2-3):313–326, 2005.
- [8] Assaf Hason and Piotr Kowalski. Strongly minimal expansions of $(\mathbb{C}, +)$ definable in o-minimal fields. (Submitted), 2006.
- [9] Assaf Hason and Alf Onshuus. Unstable structures definable in o-minimal theories. (Preprint. To appear in Proceeding of logic colloquium 2006), 2006.
- [10] Ehud Hrushovski. Locally modular regular types. In J.T. Baldwin, editor, *Classification Theory*, volume 1292 of *Lecture Notes in Mathematics*, pages 132–164. Springer-Verlag, Berlin/Heidelberg/New York, 1987.
- [11] Ehud Hrushovski. Unidimensional theories. an introduction to geometric stability theory. In H.D. Ebbinghaus et Al., editor, *Logic Colloquium '87 (Granda 1987)*, pages 73–103. North-Holland, Amsterdam, 1989.
- [12] Ehud Hrushovski. Strongly minimal expansions of algebraically closed fields. *Israel Journal of Mathematics*, 79:129–151, 1992.
- [13] Ehud Hrushovski. A new strongly minimal set. *Annals of Pure and Applied Logic*, 62:147–166, 1993.
- [14] Ehud Hrushovski and Anand Pillay. Groups definable in local fields and pseudo-finite fields. *Israel J. Math.*, 85(1-3):203–262, 1994.
- [15] Ehud Hrushovski and Boris Zilber. Zariski geometries. *Journal of the American Mathematical Society*, 9(1):1–55, 1996.
- [16] Alf Onshuus. Properties and consequences of thorn-independence. *J. Symbolic Logic*, 71(1):1–21, 2006.
- [17] Alf Onshuus and Ya'acov Peterzil. A note on stable sets and groups in theories with nip. *Preprint*, 2006.
- [18] Margarita Otero, Ya'acov Peterzil, and Anand Pillay. On groups and rings definable in o-minimal expansions of real closed fields. *Bull. London Math. Soc.*, 28(1):7–14, 1996.
- [19] Ya'acov Peterzil, Anand Pillay, and Sergei Starchenko. Simple algebraic and semialgebraic groups over real closed fields. *Trans. Amer. Math. Soc.*, 352(10):4421–4450 (electronic), 2000.
- [20] Ya'acov Peterzil and Sergei Starchenko. A trichotomy theorem for o-minimal structures. *Proc. London Math. Soc. (3)*, 77(3):481–523, 1998.
- [21] Ya'acov Peterzil and Sergei Starchenko. Expansions of algebraically closed fields in o-minimal structures. *Selecta Math. (N.S.)*, 7(3):409–445, 2001.
- [22] Anand Pillay. *Geometric stability theory*, volume 32 of *Oxford Logic Guides*. The Clarendon Press Oxford University Press, New York, 1996. Oxford Science Publications.
- [23] Bruno Poizat. *Stable groups*, volume 87 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2001. Translated from the 1987 French original by Moses Gabriel Klein.
- [24] E. D. Rabinovich. *Definability of a field in sufficiently rich incidence systems*, volume 14 of *QMW Maths Notes*. Queen Mary and Westfield College School of Mathematical Sciences, London, 1993. With an introduction by Wilfrid Hodges.
- [25] Boris Zilber. *Uncountably Categorical Theories*, volume 117. Translations of Mathematical Monographs, American Mathematical Society, Providence, RI., 1993.

UNIVERSITY OF OXFORD, MATHEMATICAL INSTITUTE, 24-29 ST GILES', OXFORD, OX1 3LB,
UK

E-mail address: `hasson@maths.ox.ac.uk`

UNIVERSIDAD DE LOS ANDES, DEPARTAMENTO DE MATEMÁTICAS, CRA. 1 No 18A-10, BO-
GOTÁ, COLOMBIA

E-mail address: `onshuus@gmail.com`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL

E-mail address: `kobi@math.haifa.ac.il`